

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

**THIS PAGE BLANK (USPTO)**

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number  
**WO 01/46165 A2**

- (51) International Patent Classification<sup>7</sup>: C07D 285/00 (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).
- (21) International Application Number: PCT/EP00/12751
- (22) International Filing Date:  
14 December 2000 (14.12.2000) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
2305/99 16 December 1999 (16.12.1999) CH (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except AT, US*): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).
- (71) Applicant (*for AT only*): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]; Brunner Strasse 59 A-1230 Vienna (AT).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): DUCRAY, Pierre [FR/FR]; 15A, rue de Huningue, F-68300 Saint-Louis (FR). BOUVIER, Jacques [CH/CH]; Rue des Parcs 2, CH-2000 Neuchatel (CH).
- Published:**  
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*


(54) Title: ORGANIC COMPOUNDS



(57) Abstract: The invention relates to compounds of general formula (I), wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub> and Q have the significances given in claim 1, and optionally the enantiomers thereof. The active ingredients have advantageous pesticidal properties. They are especially suitable for the control of pests on domestic animals and livestock.

Organic compounds

The present invention relates to new substituted aminoheterocyclamides of formula



wherein

$R_1$  is hydrogen, halogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -haloalkoxy,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -haloalkylthio,  $C_1$ - $C_6$ -alkoxy- $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkylthio- $C_1$ - $C_6$ -alkyl,  $C_3$ - $C_8$ -cycloalkyl or unsubstituted or mono- to penta-substituted phenyl, whereby the substituents are selected from the group comprising  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy, aryloxy, halogen, cyano and nitro, whereby if the number of substituents is greater than 1, the substituents may be identical or different;

$R_2$  is hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy- $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkylsulphonyl,  $C_3$ - $C_8$ -cycloalkyl,  $-COOR_6$ ,  $-CONR_7R_8$ ,  $-COR_6$ , allyl,  $-CH_2-O-R_6$ ; or heteroaryl, arylcarbonyl, arylsulphonyl or aryl- $C_1$ - $C_6$ -alkyl which are each either unsubstituted or substituted once or many times by substituents selected from the group comprising  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy, halogen, cyano, hydroxy, amino and nitro, whereby if the number of substituents is greater than 1, the substituents may be identical or different;

$R_3$  and  $R_4$ , independently of one another, are hydrogen,  $C_1$ - $C_6$ -alkyl or together with the C-atom to which they are bonded, a  $C_3$ - $C_7$ -cycloalkyl ring;

$Q$  is a single bond,  $C_1$ - $C_6$ -alkylene,  $C_2$ - $C_6$ -alkenylene,  $C_2$ - $C_6$ -alkynylene or  $C_1$ - $C_6$ -alkylenoxy;

$R_5$  is aryl or heterocyclyl, each of which is either unsubstituted or substituted once or many times by substituents selected from the group comprising  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy, aryloxy, halogen, cyano, hydroxy, amino and nitro, whereby if the number of substituents is greater than 1, the substituents may be identical or different.

$R_6$  is  $C_1$ - $C_6$ -alkyl, phenyl or benzyl;

$R_7$  and  $R_8$  independently of one another, are hydrogen or  $C_1$ - $C_6$ -alkyl;

$X_1$  and  $X_2$  independently of one another, are N, C(CN),  $-C(COOR_6)=$ ,  $-C(COR_6)=$ ,  $-C(SOR_6)=$ ,  $-C(CONR_7R_8)=$  or  $-C(NO_2)=$ ; and

$X_3$  is O or S;

whereby  $X_1$  is other than C(CN) if  $R_5$  is phenyl that is either unsubstituted or substituted once to many times;

the preparation thereof and the use thereof in the control of pests, and also pesticides containing at least one of these compounds.

Substituted aminoheterocyclamides having pesticidal activity are described for example in DE 195 42 372. However, the active ingredients specifically disclosed therein cannot always fulfil the requirements regarding potency and activity spectrum. There is therefore a need for active ingredients with improved pesticidal properties. It has now been found that the aminoheterocyclamides of formula I have excellent pesticidal properties, especially against endoparasites.

Unless otherwise defined, the general terms used hereinabove and herein below have the meanings given herein below.

Halogen - as a group *per se* and as structural element of other groups and compounds such as halogen-alkyl, halogen-alkylthio and halogen-alkoxy - is fluorine, chlorine, bromine or iodine, especially fluorine, chlorine or bromine, in particular fluorine or chlorine, especially chlorine.

If not defined to the contrary, carbon-containing groups and compounds contain 1 to 6, preferably 1 to 3, especially 1 or 2, carbon atoms.

Alkyl - as a group *per se* and as structural element of other groups and compounds such as phenylalkyl, halogen-alkyl, alkoxy, halogen-alkoxy, alkylthio and halogen-alkylthio - is, in each case with due consideration of the specific number of carbon atoms in the group or compound in question, either straight-chained, i.e. methyl, ethyl, propyl, butyl, pentyl or hexyl, or branched, e.g. isopropyl, isobutyl, sec.-butyl, tert.-butyl, isopentyl, neopentyl or isohexyl.

Cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, preferably cyclopropyl.

Alkenyl and alkynyl are straight-chained or branched and respectively contain two or preferably one unsaturated carbon-carbon bond(s). The double or triple bonds of these substituents are separated from the remaining part of compound I preferably by at least one saturated carbon atom. Allyl, methallyl, but-2-enyl, but-3-enyl, propargyl, but-2-ynyl and but-3-ynyl may be mentioned by way of example.

Halogen-substituted carbon-containing groups and compounds, such as haloalkyl, haloalkylthio, and haloalkoxy, may be partially halogenated or perhalogenated, whereby in the case of multiple halogenation, the halogen substituents may be identical or different. Examples of haloalkyl - as a group *per se* and as structural element of other groups and compounds such as haloalkylthio and haloalkoxy - are methyl which is mono- to trisubstituted by fluorine, chlorine and/or bromine, such as  $\text{CHF}_2$  or  $\text{CF}_3$ ; ethyl which is mono- to pentasubstituted by fluorine, chlorine and/or bromine, such as  $\text{CH}_2\text{CF}_3$ ,  $\text{CF}_2\text{CF}_3$ ,  $\text{CF}_2\text{CCl}_3$ ,  $\text{CF}_2\text{CHCl}_2$ ,  $\text{CF}_2\text{CHF}_2$ ,  $\text{CF}_2\text{CFCl}_2$ ,  $\text{CF}_2\text{CHBr}_2$ ,  $\text{CF}_2\text{CHClF}$ ,  $\text{CF}_2\text{CHBrF}$  or  $\text{CClFCHClF}$ ; propyl or isopropyl, mono- to heptasubstituted by fluorine, chlorine and/or bromine, such as  $\text{CH}_2\text{CHBrCH}_2\text{Br}$ ,  $\text{CF}_2\text{CHFCH}_2\text{F}$ ,  $\text{CH}_2\text{CF}_2\text{CF}_3$ ,  $\text{CF}_2\text{CF}_2\text{CF}_3$  or  $\text{CH}(\text{CF}_3)_2$ ; and butyl or one of its isomers, mono- to nonasubstituted by fluorine, chlorine and/or bromine, such as  $\text{CF}(\text{CF}_3)\text{CHFCF}_3$ ,  $\text{CF}_2(\text{CF}_2)_2\text{CF}_3$  or  $\text{CH}_2(\text{CF}_2)_2\text{CF}_3$ .

Aryl - as a group *per se* and as a structural element of other groups and compounds, such as arylcarbonyl, arylsulphonyl or aryl- $\text{C}_1\text{-C}_6$ -Alkyl, is phenyl or naphthyl, preferably phenyl.

Heteroaryl signifies an aromatic, five- or six-fold cyclic group, which contains at least one hetero atom from the group selected from oxygen, nitrogen and sulphur. Typical representatives are for example pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, pyrrol, imidazolyl, thienyl, pyrazolyl, isoxazolyl, oxazolyl and thiazolyl.

Heteroaryl signifies an aliphatic or aromatic, optionally benzo-condensed, five- or six-fold cyclic group, which contains at least one hetero atom from the group selected from oxygen, nitrogen and sulphur. Typical representatives are, for example, dioxolanyl, pyrrolidinyl, piperidyl, morpholinyl, pyridyl, pyrrol, furyl, thienyl, imidazolyl, tetrahydrofuryl, tetrahydropyrrol, tetrahydropyranyl, dihydrofuryl, dihydropyranyl, benzofuryl, benzothienyl, isoxazolyl, oxazolyl, thiazolyl, oxazolynyl, oxazolidinyl, indolyl, imidazolynyl, imidazolidinyl, dioxanyl and pyrazolyl.

Preferred embodiments in the context of the compounds of formula I are:


- (1) A compound of formula I, in which R<sub>1</sub> is halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl or phenyl; preferably halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; more preferably halogen or C<sub>1</sub>-C<sub>2</sub>-haloalkyl; most preferably chlorine or trifluoromethyl;
- (2) A compound of formula I, wherein R<sub>2</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, pyridyl or picolinyl; preferably hydrogen, C<sub>1</sub>-C<sub>2</sub>-alkyl, pyridyl or picolinyl; most preferably hydrogen, pyridyl or picolinyl;
- (3) A compound of formula I, wherein R<sub>3</sub> and R<sub>4</sub>, independently of one another, are hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, or together with the C-atom to which they are bonded, a C<sub>3</sub>-C<sub>5</sub>-cycloalkyl ring; preferably hydrogen, C<sub>1</sub>-C<sub>2</sub>-alkyl or together with the C-atom to which they are bonded, a C<sub>3</sub>-C<sub>4</sub>-cycloalkyl ring; most preferably hydrogen or together with the C-atom to which they are bonded, a cyclopropyl ring;
- (4) A compound of formula I, wherein Q is a single bond, C<sub>1</sub>-C<sub>4</sub>-alkylene, C<sub>2</sub>-C<sub>4</sub>-alkenylene or C<sub>2</sub>-C<sub>4</sub>-alkynylene; preferably a single bond or C<sub>1</sub>-C<sub>2</sub>-alkylene; most preferably a single bond or methylene;
- (5) A compound of formula I, wherein R<sub>5</sub> is aryl or heterocyclyl, each of which is unsubstituted or is substituted once or many times by substituents selected from the group comprising C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, halogen, cyano and nitro, whereby if the number of substituents is greater than 1, the substituents may be identical or different; preferably phenyl which is unsubstituted or is substituted once or many times by substituents selected from the group comprising C<sub>1</sub>-C<sub>2</sub>-alkyl, C<sub>1</sub>-C<sub>2</sub>-alkoxy and halogen, whereby if the number of substituents is greater than 1, the substituents may be identical or different; most preferably unsubstituted or mono- or disubstituted phenyl, whereby the substituents are selected from the group comprising methoxy and chlorine, whereby if the number of substituents is greater than 1, the substituents may be identical or different;
- (6) A compound of formula I, wherein X<sub>3</sub> is O;
- (7) A compound of formula I, wherein R<sub>1</sub> is halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl or phenyl; R<sub>2</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, pyridyl or picolinyl; R<sub>3</sub> and R<sub>4</sub> independently of one another, are hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or, together with the C-atom to which they are bonded, a C<sub>3</sub>-C<sub>5</sub>-cycloalkyl ring; Q is a single bond, C<sub>1</sub>-C<sub>4</sub>-alkylene, C<sub>2</sub>-C<sub>4</sub>-alkenylene or C<sub>2</sub>-C<sub>4</sub>-alkynylene; R<sub>5</sub> is aryl or heterocyclyl, each of which is unsubstituted or substituted once or many times by substituents selected from the group comprising C<sub>1</sub>-C<sub>4</sub>-

alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, halogen, cyano and nitro, whereby if the number of substituents is greater than 1, the substituents may be identical or different; and X<sub>3</sub> is O;


(8) A compound of formula I, wherein R<sub>1</sub> is halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; R<sub>2</sub> is hydrogen, C<sub>1</sub>-R<sub>2</sub>-alkyl, pyridyl or picoliny; R<sub>3</sub> and R<sub>4</sub> are hydrogen, C<sub>1</sub>-C<sub>2</sub>-alkyl or, together with the C-atom to which they are bonded, a C<sub>3</sub>-C<sub>4</sub>-cycloalkyl ring; Q is a single bond or C<sub>1</sub>-C<sub>2</sub>-alkylene; R<sub>5</sub> is phenyl which is unsubstituted or substituted once or many times by substituents selected from the group comprising C<sub>1</sub>-C<sub>2</sub>-alkyl, C<sub>1</sub>-C<sub>2</sub>-alkoxy and halogen, whereby if the number of substituents is greater than 1, the substituents may be identical or different; and X<sub>3</sub> is O;

(9) A compound of formula I, wherein R<sub>1</sub> is chlorine or trifluoromethyl; R<sub>2</sub> is hydrogen, pyridyl or picoliny, R<sub>3</sub> and R<sub>4</sub> are hydrogen, or together with the C-atom to which they are bonded, a cyclopropyl ring; Q is a single bond or methylene; R<sub>5</sub> is unsubstituted or mono- or disubstituted phenyl, whereby the substituents are selected from the group comprising methoxy and chlorine, whereby if the number of substituents is greater than 1, the substituents may be identical or different; and X<sub>3</sub> is O.

A further object of the invention is the process for the preparation of the compounds of formula I and optionally the enantiomers thereof, for example characterised in that a compound of formula



which is known or may be produced analogously to corresponding known compounds, and wherein R<sub>1</sub>, R<sub>2</sub>, X<sub>1</sub> and X<sub>2</sub> are defined as given for formula I, is reacted with a compound of formula



which is known or may be produced analogously to corresponding known compounds, in which X<sub>3</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and Q are defined as for formula I, and Z is a leaving group, optionally



in the presence of a basic catalyst, and if desired, a compound of formula I which is obtainable by this process or in another way, or an enantiomer thereof, may be converted into another compound of formula I or an enantiomer thereof, a mixture of enantiomers which is obtainable by this process is separated and the desired enantiomer isolated.

Suitable leaving groups are halogen, C<sub>1</sub>-C<sub>6</sub>-alkoxy or hydroxy, preferably chlorine.

Suitable bases for facilitating the reaction are e.g. trialkylamines, basic heterocycles or phosphines. Triethylamine, diisopropylethylamine, pyridine, 4-(N,N-dimethylamino)pyridine, quinuclidine, 1,5-diazabicyclo[5.4.0]undec-5-ene (DBU) and triphenylphosphine may be mentioned by way of example. Diisopropylethylamine is preferred.

The reaction partners can be reacted with one another as they are, i.e. without the addition of a solvent or diluent, e.g. in the melt. In most cases, however, the addition of an inert solvent or diluent, or a mixture thereof, is of advantage. Examples of such solvents or diluents are: aromatic, aliphatic and alicyclic hydrocarbons and halogenated hydrocarbons, such as benzene, toluene, xylene, mesitylene, tetraline, chlorobenzene, dichlorobenzene, bromobenzene, petroleum ether, hexane, cyclohexane, dichloromethane, trichloromethane, tetrachloromethane, dichloroethane, trichloroethene or tetrachloroethene; ethers, such as diethyl ether, dipropyl ether, diisopropyl ether, dibutyl ether, tert-butyl methyl ether, ethylene glycol monomethyl ether, ethylene glycol monoethyl ether, ethylene glycol dimethylether, dimethoxydiethylether, tetrahydrofuran or dioxane; ketones such as acetone, methyl ethyl ketone or methyl isobutyl ketone; amides such as N,N-dimethylformamide, N,N-diethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone or hexamethylphosphoric acid triamide; nitriles such as acetonitrile or propionitrile; and sulphoxides, such as dimethyl sulphoxide. If the reaction takes place in the presence of a base, then bases used in excess, such as triethylamine, pyridine, N-methylmorpholine, or N,N-diethylaniline, can also serve as solvents or diluents. Preferably, halogenated hydrocarbons are used, especially dichloromethane.

The reaction is advantageously carried out in a temperature range of ca. -20°C to ca. +150°C, preferably from ca. -10°C to ca. +80°C, most preferably from ca. 0°C to ca. +40°C.

In a preferred embodiment, a compound of formula II is reacted at 0° to 120°, preferably 20°, in a halogenated hydrocarbon, preferably dichloromethane, with a compound of formula III.

The compounds I may be present in the form of one of the possible isomers or as a mixture thereof, e.g. depending on the number, absolute and relative configurations of the asymmetric carbon atoms as pure isomers, such as antipodes and/or diastereoisomers, or as isomeric mixtures, such as enantiomeric mixtures, e.g. racemates, diastereoisomeric mixtures or racemic mixtures; the invention relates to both the pure isomers and all the possible isomeric mixtures, and is to be understood as such hereinbefore and hereinafter, even if stereochemical details are not specifically mentioned in each case.

Depending on the choice of starting materials and procedures, diastereoisomeric mixtures and racemic mixtures of compounds I, which are obtained in accordance with the invention or in another way, may be separated in known manner into the pure diastereoisomers or racemates based on the physical-chemical differences of the constituents, for example by means of fractional crystallisation, distillation and/or chromatography.

Mixtures of enantiomers that are obtainable accordingly, such as racemates, may be broken down into the optical antipodes by known methods, for example by recrystallisation from an optically active solvent, by chromatography on chiral adsorbents, e.g. high-pressure liquid chromatography (HPLC) on acetyl cellulose, with the assistance of appropriate micro-organisms, by cleavage with specific immobilised enzymes, through the formation of inclusion compounds, e.g. using chiral crown ethers, wherein only one enantiomer is complexed.

According to the invention, apart from isolation of corresponding isomer mixtures, generally known methods of diastereoselective or enantioselective synthesis can also be applied to obtain pure diastereoisomers or enantiomers, e.g. by carrying out the method of the invention using educts with correspondingly suitable stereochemistry.

It is advantageous to isolate or synthesize the biologically more active isomer in each case, e.g. enantiomer or isomer mixture, e.g. enantiomer mixture, if the individual components show differences in biological efficacy.

In the method of the present invention, the starting materials and intermediates used are preferably those that lead to the compounds I described at the beginning as being especially useful.

The invention relates especially to the method of preparation described in the example.

Starting materials and intermediates, which are new and are used according to the invention for the preparation of compounds I, as well as their usage and process for the preparation thereof, similarly form an object of the invention.

The compounds I according to the invention are notable for their broad activity spectrum and are valuable active ingredients for use in pest control, including in particular the control of endo- and ecto-parasites on animals, whilst being well-tolerated by warm-blooded animals, fish and plants,

In the context of the present invention, ectoparasites are understood to be in particular insects, mites and ticks. These include insects of the order: *Lepidoptera*, *Coleoptera*, *Homoptera*, *Heteroptera*, *Diptera*, *Thysanoptera*, *Orthoptera*, *Anoplura*, *Siphonaptera*, *Mallophaga*, *Thysanura*, *Isoptera*, *Psocoptera* and *Hymenoptera*. However, the ectoparasites which may be mentioned in particular are those which trouble humans or animals and carry pathogens, for example flies such as *Musca domestica*, *Musca vetustissima*, *Musca autumnalis*, *Fannia canicularis*, *Sarcophaga carnaria*, *Lucilia cuprina*, *Hypoderma bovis*, *Hypoderma lineatum*, *Chrysomya chloropyga*, *Dermatobia hominis*, *Cochliomyia hominivorax*, *Gasterophilus intestinalis*, *Oestrus ovis*, *Stomoxys calcitrans*, *Haematobia irritans* and midges (*Nematocera*), such as *Culicidae*, *Simuliidae*, *Psychodidae*, but also blood-sucking parasites, for example fleas, such as *Ctenocephalides felis* and *Ctenocephalides canis* (cat and dog fleas), *Xenopsylla cheopis*, *Pulex irritans*, *Dermatophilus penetrans*, lice, such as *Damalina ovis*, *Pediculus humanis*, biting flies and horse-flies (*Tabanidae*), *Haematopota* spp. such as *Haematopota pluvialis*, *Tabanidea* spp. such as *Tabanus nigrovittatus*, *Chrysopsinae* spp. such as *Chrysops caecutiens*, tsetse flies, such as species of *Glossinia*, biting insects, particularly cockroaches, such as *Blatella germanica*, *Blatta orientalis*, *Periplaneta americana*, mites, such as *Dermanyssus gallinae*, *Sarcoptes scabiei*, *Psoroptes ovis* and *Psorergates* spp. and last but not least ticks. The latter belong to the order *Acarina*. Known representatives of ticks are, for example, *Boophilus*, *Amblyomma*, *Anocentor*, *Dermacentor*, *Haemaphysalis*, *Hyalomma*, *Ixodes*, *Rhipicentor*, *Margaropus*, *Rhipicephalus*, *Argas*, *Otobius* and *Ornithodoros* and the like, which preferably infest warm-blooded animals including farm animals, such as cattle, pigs, sheep and goats, poultry such as chickens, turkeys and geese, fur-bearing animals such as mink, foxes, chinchillas, rabbits and the like, as well as domestic animals such as cats and dogs, but also humans.

Compounds I can also be used against hygiene pests, especially of the order *Diptera* of the families *Sarcophagidae*, *Anophilidae* and *Culicidae*; the orders *Orthoptera*, *Dictyoptera* (e.g. the family *Blattidae*) and *Hymenoptera* (e.g. the family *Formicidae*).

Compounds I also have sustainable efficacy on parasitic mites and insects of plants. In the case of spider mites of the order *Acarina*, they are effective against eggs, nymphs and adults of *Tetranychidae* (*Tetranychus* spp. and *Panonychus* spp.).

They have high activity against sucking insects of the order *Homoptera*, especially against pests of the families *Aphididae*, *Delphacidae*, *Cicadellidae*, *Psyllidae*, *Loccidae*, *Diaspididae* and *Eriophyidae* (e.g. rust mite on citrus fruits); the orders *Hemiptera*, *Heteroptera* and *Thysanoptera*, and on the plant-eating insects of the orders *Lepidoptera*, *Coleoptera*, *Diptera* and *Orthoptera*.

They are similarly suitable as a soil insecticide against pests in the soil.

The compounds of formula I are therefore effective against all stages of development of sucking insects and eating insects on crops such as cereals, cotton, rice, maize, soya, potatoes, vegetables, fruit, tobacco, hops, citrus, avocados and other crops.

The compounds of formula I are also effective against plant nematodes of the species *Meloidogyne*, *Heterodera*, *Pratylenchus*, *Ditylenchus*, *Radopholus*, *Rizoglyphus* etc.

In particular, the compounds are effective against helminths, in which the endoparasitic nematodes may be the cause of serious diseases of mammals and poultry, e.g. sheep, pigs, goats, cattle, horses, donkeys, dogs, cats, guinea-pigs and exotic birds. Typical nematodes of this indication are: *Haemonchus*, *Trichostrongylus*, *Ostertagia*, *Nematodirus*, *Cooperia*, *Ascaris*, *Bunostomum*, *Oesophagostomum*, *Charbertia*, *Trichuris*, *Strongylus*, *Trichonema*, *Dictyocaulus*, *Capillaria*, *Heterakis*, *Toxocara*, *Ascaridia*, *Oxyuris*, *Ancylostoma*, *Uncinaria*, *Toxascaris* and *Parascaris*. The particular advantage of the compounds of formula I is their efficacy against those parasites that are resistant towards active ingredients based on benzimidazole.

Certain pests of the species *Nematodirus*, *Cooperia* and *Oesophagostomum* infest the intestinal tract of the host animal, while others of the species *Haemonchus* and *Ostertagia* are parasitic in the stomach and those of the species *Dictyocaulus* are parasitic in the lung tissue. Parasites of the families *Filariidae* and *Setariidae* may be found in the internal cell tissue and in the organs, e.g. the heart, the blood vessels, the lymph vessels and the

subcutaneous tissue. A particularly notable parasite is the heartworm of the dog, *Dirofilaria immitis*. The compounds of formula I are highly effective against these parasites.

Furthermore, the compounds of formula I are also especially suitable for the control of human pathogenic parasites. Of these, typical representatives that appear in the digestive tract are those of the species *Ancylostoma*, *Necator*, *Ascaris*, *Strongyloides*, *Trichinella*, *Capillaria*, *Trichuris* and *Enterobius*. The compounds of the present invention are also effective against parasites of the species *Wuchereria*, *Brugia*, *Onchocerca* and *Loa* from the family of *Filariidae*, which appear in the blood, in the tissue and in various organs, and also against *Dracunculus* and parasites of the species *Strongyloides* and *Trichinella*, which infect the gastrointestinal tract in particular.

The good pesticidal activity of the compounds of formula I corresponds to a mortality rate of at least 50-60% of the pests mentioned. In particular, the compounds of formula I are notable for the exceptionally long duration of efficacy.

The activity of the compounds according to the invention and of the compositions containing them against animal pests may be substantially broadened and adapted to the prevailing circumstances by adding other insecticides and/or acaricides. The additives in question may be for example representatives of the following classes of active ingredient: organophosphorus compounds, nitrophenols and derivatives, formamidines, ureas, carbamates, pyrethroids, chlorinated hydrocarbons, neonicotinoids and *Bacillus thuringiensis* preparations.

The compounds of formula I are preferably employed in unmodified form or preferably together with the adjuvants conventionally used in the art of formulation and may therefore be processed in a known manner to give, for example, emulsifiable concentrates, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granules or microencapsulations in polymeric substances. As with the compositions, the methods of application, such as spraying, atomising, dusting, scattering or pouring, are selected in accordance with the intended objectives and the prevailing circumstances.

The formulation, i.e. the agents, preparations or compositions containing the active ingredient of formula I, or combinations of these active ingredients with other agrochemical active ingredients, and optionally a solid or liquid adjuvant, are produced in a manner known *per se*, for example by intimately mixing and/or grinding the active ingredients with

spreading compositions, for example with solvents, solid carriers, and optionally surface-active compounds (surfactants).

The solvents in question may be: aromatic hydrocarbons, preferably fractions of alkylbenzenes having 8 to 12 carbon atoms, such as xylene mixtures or alkylated naphthalenes, aliphatic or cyclo-aliphatic hydrocarbons, such as cyclohexane, paraffins or tetrahydronaphthalene, alcohols, such as ethanol, propanol or butanol, glycols and their ethers and esters, such as propylene glycol, dipropylene glycol ether, ethylene glycol or ethylene glycol monomethyl or -ethyl ether, ketones, such as cyclohexanone, isophorone or diacetanol alcohol, strong polar solvents, such as N-methyl-2-pyrrolidone, dimethyl sulphoxide or dimethylformamide, or water, vegetable oils, such as rape, castor, coconut, or soybean oil, and also, if appropriate, silicone oils.

The solid carriers used for example for dusts and dispersible powders, are normally natural mineral fillers such as calcite, talcum, kaolin, montmorillonite or attapulgit. In order to improve the physical properties it is also possible to add highly dispersed silicic acid or highly dispersed absorbent polymers. Suitable granulated adsorptive carriers are porous types, for example pumice, broken brick, sepiolite or bentonite, and suitable non-sorbent carriers are materials such as calcite or sand. In addition, a great number of pregranulated materials of inorganic or organic nature can be used, e.g. especially dolomite or pulverised plant residues.

Depending on the type of active ingredient of formula I to be formulated, or the combination of these active ingredients with other insecticides or acaricides, suitable surface-active compounds are non-ionic, cationic and/or anionic surfactants having good emulsifying, dispersing and wetting properties. The surfactants are also understood to be surfactant mixtures.

Suitable anionic surfactants can be both so-called water-soluble soaps and water-soluble synthetic surfactant compounds.

Suitable soaps are the alkali metal salts, alkaline earth metal salts or unsubstituted or substituted ammonium salts of higher fatty acids ( $C_{10}-C_{22}$ ), for example the sodium or potassium salts of oleic or stearic acid, or of natural fatty acid mixtures which can be obtained for example from coconut oil or tallow oil. The fatty acid methyltaurine salts may also be mentioned as surfactants.

More frequently, however, so-called synthetic surfactants are used, especially fatty sulphonates, fatty sulphates, sulphonated benzimidazole derivatives or alkylarylsulphonates.

The fatty sulphonates or sulphates are usually in the form of alkali metal salts, alkaline earth metal salts or unsubstituted or substituted ammoniums salts and have an alkyl radical with 8 to 22 carbon atoms, which also includes the alkyl moiety of acyl radicals, for example, the sodium or calcium salt of ligninsulphonic acid, of dodecylsulphate or of a mixture of fatty alcohol sulphates obtained from natural fatty acids. These compounds also comprise the salts of sulphuric acid esters and sulphonic acids of fatty alcohol/ethylene oxide adducts. The sulphonated benzimidazole derivatives preferably contain 2 sulphonic acid groups and one fatty acid radical containing 8 to 22 carbon atoms. Examples of alkylarylsulphonates are the sodium, calcium or triethanolamine salts of dodecylbenzenesulphonic acid, dibutyl-naphthalenesulphonic acid, or of a naphthalenesulphonic acid / formaldehyde condensation product. Also suitable are corresponding phosphates, e.g. salts of the phosphoric acid ester of an adduct of p-nonylphenol with 4 to 14 moles of ethylene oxide or phospholipids.

Non-ionic surfactants are preferably polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols, or saturated or unsaturated fatty acids and alkylphenols, said derivatives containing 3 to 30 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon moiety and 6 to 18 carbon atoms in the alkyl moiety of the alkylphenols. Further suitable non-ionic surfactants are the water-soluble adducts of polyethylene oxide with polypropylene glycol, ethylenediamine polypropylene glycol and alkylpolypropylene glycol containing 1 to 10 carbon atoms in the alkyl chain, which adducts contain 20 to 250 ethylene glycol ether groups and 10 to 100 propylene glycol ether groups. These compounds usually contain 1 to 5 ethylene glycol units per propylene glycol unit.

Examples of non-ionic surfactants are nonylphenolpolyethoxyethanols, castor oil polyglycol ethers, polypropylene/polyethylene oxide adducts, tributylphenoxypolyethoxyethanol, polyethylene glycol and octylphenoxypolyethoxyethanol. Also suitable are fatty acid esters of polyoxyethylene sorbitan, such as polyoxyethylene sorbitan trioleate.

Cationic surfactants are preferably quaternary ammonium salts which have as N-substituent at least one C<sub>8</sub>-C<sub>22</sub> alkyl radical and, as further substituents, lower - where appropriate - halogenated alkyl, benzyl or lower hydroxyalkyl radicals. The salts preferably exist as

halides, methyl sulphates or ethyl sulphates, preferably as stearyl trimethylammonium chloride or benzyl-di-(2-chloroethyl)-ethylammonium bromide.

The surfactants which are customary in formulation technology are described for example in the following publications:

"Mc Cutcheon's Detergents and Emulsifiers Annual", McPublishing Corp., Glen Rock, NJ, USA, 1988",

H. Stache, "Tensid-Taschenbuch" (Surfactants Handbook), 2<sup>nd</sup> edition, C. Hanser Publishing Munich, Vienna 1981.

M. and J. Ash. "Encyclopaedia of Surfactants", Vol. I-III, Chemical Publishing Co., New York, 1980-1981.

Preferred application forms for usage on warm-blooded animals in the control of helminths include solutions, emulsions, suspensions (drenches), food additives, powders, tablets including effervescent tablets, boli, capsules, micro-capsules and pour-on formulations, whereby the physiological compatibility of the formulation excipients must be taken into consideration.

The binders for tablets and boli may be chemically modified polymeric natural substances that are soluble in water or in alcohol, such as starch, cellulose or protein derivatives (e.g. methyl cellulose, carboxymethyl cellulose, ethylhydroxyethyl cellulose, proteins such as zein, gelatin and the like), as well as synthetic polymers, such as polyvinyl alcohol, polyvinyl pyrrolidone etc. The tablets also contain fillers (e.g. starch, microcrystalline cellulose, sugar, lactose etc.), glidants and disintegrants.

If the anthelmintics are present in the form of feed concentrates, then the carriers used are e.g. performance feeds, feed grain or protein concentrates. Such feed concentrates or compositions may contain, apart from the active ingredients, also additives, vitamins, antibiotics, chemotherapeutics or other pesticides, primarily bacteriostats, fungistats, coccidiostats, or even hormone preparations, substances having anabolic action or substances which promote growth, which affect the quality of meat of animals for slaughter or which are beneficial to the organism in another way. If the compositions or the active ingredients of formula I contained therein are added directly to feed or to the drinking troughs, then the formulated feed or drink contains the active ingredients preferably in a concentration of ca. 0.0005 to 0.02 % by weight (5-200 ppm).



Application of the compositions according to the invention to the animals to be treated may take place topically, perorally, parenterally or subcutaneously, the composition being present in the form of solutions, emulsions, suspensions, (drenches), powders, tablets, boli, capsules and pour-on formulations.

The compounds of formula I according to the invention may be used alone or in combination with other biocides. They may be combined with pesticides having the same sphere of activity e.g. to increase activity, or with substances having another sphere of activity e.g. to broaden the range of activity. It can also be sensible to add so-called repellents. If the range of activity is to be extended to endoparasites, e.g. wormers, the compounds of formula I are suitably combined with substances having endoparasitic properties. Of course, they can also be used in combination with antibacterial compositions. Since the compounds of formula I are adulticides, i.e. since they are effective in particular against the adult stage of the target parasites, the addition of pesticides which instead attack the juvenile stages of the parasites may be very advantageous. In this way, the greatest part of those parasites that produce great economic damage will be covered. Moreover, this action will contribute substantially to avoiding the formation of resistance. Many combinations may also lead to synergistic effects, i.e. the total amount of active ingredient can be reduced, which is desirable from an ecological point of view. Preferred groups of combination partners and especially preferred combination partners are named in the following, whereby combinations may contain one or more of these partners in addition to a compound of formula I.

Suitable partners in the mixture may be biocides, e.g. the insecticides and acaricides with a varying mechanism of activity, which are named in the following and have been known to the person skilled in the art for a long time, e.g. chitin synthesis inhibitors, growth regulators; active ingredients which act as juvenile hormones; active ingredients which act as adulticides; broad-band insecticides, broad-band acaricides and nematocides; and also the well known anthelmintics and insect- and/or acarid-deterring substances, said repellents or detachers.

Non-limitative examples of suitable insecticides and acaricides are:

1. Abamectin
2. AC 303 630
3. Acephat
4. Acrinathrin

5. Alanycarb
6. Aldicarb
7. $\alpha$ -Cypermethrin
8. Alphamethrin

9. Amitraz
10. Avermectin B <sub>1</sub>
11. AZ 60541
12. Azinphos A

13. Azinphos M
14. Azinphos-methyl
15. Azocyclotin
16. <i>Bacillus subtil.</i> toxin
17. Bendiocarb
18. Benfuracarb
19. Bensultap
20. $\beta$ -Cyfluthrin
21. Bifenthrin
22. BPMC
23. Brofenprox
24. Bromophos A
25. Bufencarb
26. Buprofezin
27. Butocarboxin
28. Butylpyridaben
29. Cadusafos
30. Carbaryl
31. Carbofuran
32. Carbophenthion
33. Cartap
34. Chloethocarb
35. Chlorethoxyfos
36. Chlorfenapyr
37. Chlorfluazuron
38. Chlormephos
39. Chlorpyrifos
40. Cis-Resmethrin
41. Clocythrin
42. Clofentezin
43. Cyanophos
44. Cycloprothrin

45. Cyfluthrin
46. Cyhexatin
47. D 2341
48. Deltamethrin
49. Demeton M
50. Demeton S
51. Demeton-S-methyl
52. Dibutylaminothio
53. Dichlofenthion
54. Dicliphos
55. Diethion
56. Diflubenzuron
57. Dimethoat
58. Dimethylvinphos
59. Dioxathion
60. DPX-MP062
61. Edifenphos
62. Enamectin
63. Endosulfan
64. Esfenvalerat
65. Ethiofencarb
66. Ethion
67. Ethofenprox
68. Ethoprophos
69. Etrimphos
70. Fenamiphos
71. Fenazaquin
72. Fenbutatinoxid
73. Fenitrothion
74. Fenobucarb
75. Fenobucarb
76. Fenothiocarb

77. Fenoxycarb
78. Fenpropathrin
79. Fenpyrad
80. Fenpyroximate
81. Fenthion
82. Fenvalerate
83. Fipronil
84. Fluazinam
85. Fluazuron
86. Flucycloxuron
87. Flucythrinate
88. Flufenoxuron
89. Flufenprox
90. Fonophos
91. Formothion
92. Fosthiazat
93. Fubfenprox
94. HCH
95. Heptenophos
96. Hexaflumuron
97. Hexythiazox
98. Hydroprene
99. Imidacloprid
100. Insect-active fungi
101. Insect-active nematodes
102. Insect-active viruses
103. Iprobenfos
104. Isofenphos
105. Isoprocarb
106. Isoxathion
107. Ivermectin

108. $\lambda$ -Cyhalothrin	136. Phosmet	164. Teflubenzuron
109. Lufenuron	137. Phoxim	165. Tefluthrin
110. Malathion	138. Pirimicarb	166. Temephos
111. Mecarbam	139. Pirimiphos A	167. Terbam
112. Mesulfenphos	140. Pirimiphos M	168. Terbufos
113. Metaldehyd	141. Promecarb	169. Tetrachlorvinphos
114. Methamidophos	142. Propaphos	170. Thiafenox
115. Methiocarb	143. Propoxur	171. Thiodicarb
116. Methomyl	144. Prothiofos	172. Thiofanox
117. Methoprene	145. Prothoat	173. Thionazin
118. Metolcarb	146. Pyrachlophos	174. Thuringiensin
119. Mevinphos	147. Pyradaphenthion	175. Tralomethrin
120. Milbemectin	148. Pyresmethrin	176. Triarthen
121. Moxidectin	149. Pyrethrum	177. Triazamate
122. Naled	150. Pyridaben	178. Triazophos
123. NC 184	151. Pyrimidifen	179. Triazuron
124. NI-25, Acetamiprid	152. Pyriproxyfen	180. Trichlorfon
125. Nitenpyram	153. RH 5992	181. Triflumuron
126. Omethoat	154. RH-2485	182. Trimethacarb
127. Oxamyl	155. Salithion	183. Vamidothion
128. Oxydemethon M	156. Sebufos	184. XMC (3,5,-Xylyl-methylcarbamate)
129. Oxydeprofos	157. Silafluofen	185. Xylylcarb
130. Parathion	158. Spinosad	186. YI 5301/5302
131. Parathion-methyl	159. Sulfotep	187. $\zeta$ -Cypermethrin
132. Permethrin	160. Sulprofos	188. Zetamethrin
133. Phenthoat	161. Tebufenozide	
134. Phorat	162. Tebufenpyrad	
135. Phosalone	163. Tebupirimphos	

Non-limitative examples of suitable anthelmintics are named in the following, a few representatives have insecticidal and acaricidal activity in addition to the anthelmintic activity, and are partly already in the above list.

- (A1) Praziquantel = 2-cyclohexylcarbonyl-4-oxo-1,2,3,6,7,11b-hexahydro-4H-pyrazino[2,1- $\alpha$ ]isoquinoline
- (A2) Closantel = 3,5-diiodo-N-[5-chloro-2-methyl-4-( $\alpha$ -cyano-4-chlorobenzyl)phenyl]-salicylamide
- (A3) Triclabendazole = 5-chloro-6-(2,3-dichlorophenoxy)-2-methylthio-1H-benzimidazole
- (A4) Levamisole = L-(-)-2,3,5,6-tetrahydro-6-phenylimidazo[2,1b]thiazole
- (A5) Mebendazole = (5-benzoyl-1H-benzimidazol-2-yl)carbaminic acid methyl ester
- (A6) Omphalotin = a macrocyclic fermentation product of the fungus *Omphalotus olearius* described in WO 97/20857
- (A7) Abamectin = avermectin B1
- (A8) Ivermectin = 22,23-dihydroavermectin B1
- (A9) Moxidectin = 5-O-demethyl-28-deoxy-25-(1,3-dimethyl-1-butenyl)-6,28- epoxy-23-(methoxyimino)-milbemycin B
- (A10) Doramectin = 25-cyclohexyl-5-O-demethyl-25-de(1-methylpropyl)-avermectin A1a
- (A11) Milbemectin = mixture of milbemycin A3 and milbemycin A4
- (A12) Milbemycinoxim = 5-oxim of milbemectin

Non-limitative examples of suitable repellents and detachers are:

- (R1) DEET (N, N-diethyl-m-toluamide)
- (R2) KBR 3023 N-butyl-2-oxycarbonyl-(2-hydroxy)-piperidine
- (R3) Cymiazole = N,-2,3-dihydro-3-methyl-1,3-thiazol-2-ylidene-2,4-xylidene

The said partners in the mixture are best known to specialists in this field. Most are described in various editions of the Pesticide Manual, The British Crop Protection Council, London, and others in the various editions of The Merck Index, Merck & Co., Inc., Rahway, New Jersey, USA or in patent literature. Therefore, the following listing is restricted to a few places where they may be found by way of example.

- (I) 2-Methyl-2-(methylthio)propionaldehyd-O-Methylcarbamoxyloxime (Aldicarb), from The Pesticide Manual, 11<sup>th</sup> Ed. (1997), The British Crop Protection Council, London, page 26;
- (II) S-(3,4-Dihydro-4-oxobenzo[d]-[1,2,3]-triazin-3-ylmethyl)O,O-dimethyl-phosphorodithioate (Azinphos-methyl), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 67;
- (III) Ethyl-N-[2,3-dihydro-2,2-dimethylbenzofuran-7-yloxycarbonyl-(methyl)aminothio]-N-isopropyl- $\beta$ -alaninate (Benfuracarb), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 96;

- (IV) 2-Methylbiphenyl-3-ylmethyl-(Z)-(1*RS*)-*cis*-3-(2-chloro-3,3,3-trifluoroprop-1-enyl)-2,2-dimethylcyclopropanecarboxylate (Bifenthrin), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 118;
- (V) 2-tert-Butylimino-3-isopropyl-5-phenyl-1,3,5-thiadiazian-4-one (Buprofezin), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 157;
- (VI) 2,3-Dihydro-2,2-dimethylbenzofuran-7-yl-methylcarbamate (Carbofuran), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 186;
- (VII) 2,3-Dihydro-2,2-dimethylbenzofuran-7-yl-(dibutylaminothio)methylcarbamate (Carbosulfan), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 188;
- (VIII) *S,S'*-(2-Dimethylaminotrimethylene)-bis(thiocarbamate) (Cartap), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 193;
- (IX) 1-[3,5-Dichloro-4-(3-chloro-5-trifluoromethyl-2-pyridyloxy)phenyl]-3-(2,6-difluorobenzoyl)-urea (Chlorfluazuron), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 213;
- (X) *O,O*-Diethyl-*O*-3,5,6-trichloro-2-pyridyl-phosphorothioate (Chlorpyrifos), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 235;
- (XI) (*RS*)- $\alpha$ -Cyano-4-fluoro-3-phenoxybenzyl-(1*RS*,3*RS*;1*RS*,3*RS*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (Cyfluthrin), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 293;
- (XII) Mixture of (*S*)- $\alpha$ -Cyano-3-phenoxybenzyl-(Z)-(1*R*,3*R*)-3-(2-chloro-3,3,3-trifluoropropenyl)-2,2-dimethylcyclopropanecarboxylate and (*R*)- $\alpha$ -cyano-3-phenoxybenzyl-(Z)-(1*R*,3*R*)-3-(2-chloro-3,3,3-trifluoropropenyl)-2,2-dimethylcyclopropanecarboxylate (Lambda-Cyhalothrin), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 300;
- (XIII) Racemate consisting of (*S*)- $\alpha$ -cyano-3-phenoxybenzyl-(1*R*,3*R*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (*R*)- $\alpha$ -cyano-3-phenoxybenzyl-(1*S*,3*S*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (Alpha-cypermethrin), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 308;

- (XIV) Mixture of the stereoisomers of (*S*)- $\alpha$ -cyano-3-phenoxybenzyl (1*RS*,3*RS*,1*RS*,3*RS*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (zeta-Cypermethrin), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 314;
- (XV) (*S*)- $\alpha$ -cyano-3-phenoxybenzyl-(1*R*,3*R*)-3-(2,2-dibromovinyl)-2,2-dimethylcyclopropanecarboxylate (Deltamethrin), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 344;
- (XVI) (4-Chlorophenyl)-3-(2,6-difluorobenzoyl)urea (Diflubenzuron), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 395;
- (XVII) (1,4,5,6,7,7-Hexachloro-8,9,10-trinorborn-5-en-2,3-ylenebismethylene)-sulphite (Endosulfan), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 459;
- (XVIII)  $\alpha$ -Ethylthio-*o*-tolyl-methylcarbamate (Ethiofencarb), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 479;
- (XIX) *O,O*-Dimethyl-*O*-4-nitro-*m*-tolyl-phosphorothioate (Fenitrothion), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 514;
- (XX) 2-*sec*-Butylphenyl-methylcarbamate (Fenobucarb), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 516;
- (XXI) (*RS*)- $\alpha$ -Cyano-3-phenoxybenzyl-(*RS*)-2-(4-chlorophenyl)-3-methylbutyrate (Fenvalerate), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 539;
- (XXII) *S*-[Formyl(methyl)carbamoylmethyl]-*O,O*-dimethyl-phosphorodithioate (Formothion), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 625;
- (XXIII) 4-Methylthio-3,5-xylyl-methylcarbamate (Methiocarb), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 813;
- (XXIV) 7-Chlorobicyclo[3.2.0]hepta-2,6-dien-6-yl-dimethylphosphate (Heptenophos), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 670;
- (XXV) 1-(6-Chloro-3-pyridylmethyl)-*N*-nitroimidazolidin-2-ylidenamine (Imidacloprid), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 706;

- (XXVI) 2-Isopropylphenyl-methylcarbamate (Isoprocarb), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 729;
- (XXVII) *O,S*-Dimethyl-phosphoramidothioate (Methamidophos), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 808;
- (XXVIII) *S*-Methyl-*N*-(methylcarbamoyloxy)thioacetimidate (Methomyl), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 815;
- (XXIX) Methyl-3-(dimethoxyphosphinyloxy)but-2-enoate (Mevinphos), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 844;
- (XXX) *O,O*-Diethyl-*O*-4-nitrophenyl-phosphorothioate (Parathion), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 926;
- (XXXI) *O,O*-Dimethyl-*O*-4-nitrophenyl-phosphorothioate (Parathion-methyl), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 928;
- (XXXII) *S*-6-Chloro-2,3-dihydro-2-oxo-1,3-benzoxazol-3-ylmethyl-*O,O*-diethyl-phosphorodithioate (Phosalone), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 963;
- (XXXIII) 2-Dimethylamino-5,6-dimethylpyrimidin-4-yl-dimethylcarbamate (Pirimicarb), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 985;
- (XXXIV) 2-Isopropoxyphenyl-methylcarbamate (Propoxur), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1036;
- (XXXV) 1-(3,5-Dichloro-2,4-difluorophenyl)-3-(2,6-difluorobenzoyl)urea (Teflubenzuron), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1158;
- (XXXVI) *S*-tert-butylthiomethyl-*O,O*-dimethyl-phosphorodithioate (Terbufos), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1165;
- (XXXVII) Ethyl-(3-*tert*.-butyl-1-dimethylcarbamoyl-1*H*-1,2,4-triazol-5-yl-thio)-acetate, (Triazamate), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1224;
- (XXXVIII) Abamectin, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 3;

- (XXXIX) 2-sec-butylphenyl-methylcarbamate (Fenobucarb), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 516;
- (XL) *N*-tert.-butyl-*N*-(4-ethylbenzoyl)-3,5-dimethylbenzohydrazide (Tebufenozide), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1147;
- (XLI) (±)-5-Amino-1-(2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-*p*-tolyl)-4-trifluoromethyl-sulphinylpyrazol-3-carbonitrile (Fipronil), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 545;
- (XLII) (*RS*)- $\alpha$ -cyano-4-fluoro-3-phenoxybenzyl(1*RS*,3*RS*;1*RS*,3*RS*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (beta-Cyfluthrin), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 295;
- (XLIII) (4-Ethoxyphenyl)-[3-(4-fluoro-3-phenoxyphenyl)propyl](dimethyl)silane (Silaf luofen), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1105;
- (XLIV) tert.-butyl (*E*)- $\alpha$ -(1,3-dimethyl-5-phenoxy-pyrazol-4-yl-methylenamino-oxy)-*p*-toluate (Fenpyroximate), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 530;
- (XLV) 2-tert.-butyl-5-(4-tert.-butylbenzylthio)-4-chloropyridazin-3(2*H*)-one (Pyridaben), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1161;
- (XLVI) 4-[[4-(1,1-dimethylphenyl)phenyl]ethoxy]-quinazoline (Fenazaquin), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 507;
- (XLVII) 4-Phenoxyphenyl-(*RS*)-2-(pyridyloxy)propyl-ether (Pyriproxyfen), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1073;
- (XLVIII) 5-Chloro-*N*-(2-[4-(2-ethoxyethyl)-2,3-dimethylphenoxy]ethyl)-6-ethylpyrimidin-4-amine (Pyrimidifen), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1070;
- (XLIX) (*E*)-*N*-(6-chloro-3-pyridylmethyl)-*N*-ethyl-*N*-methyl-2-nitrovinylidenediamine (Nitenpyram), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 880;



- (L) (E)-N<sup>1</sup>-[(6-chloro-3-pyridyl)methyl]-N<sup>2</sup>-cyano-N<sup>1</sup>-methylacetamidine (NI-25, Acetamiprid), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 9;
- (LI) Avermectin B<sub>1</sub>, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 3;
- (LII) an insect-active extract from a plant, especially (2R,6aS,12aS)-1,2,6,6a,12,12a-hexhydro-2-isopropenyl-8,9-dimethoxy-chromeno[3,4-b]furo[2,3-h]chromen-6-one (Rotenone), from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1097; and an extract from *Azadirachta indica*, especially azadirachtin, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 59; and
- (LIII) a preparation which contains insect-active nematodes, preferably *Heterorhabditis bacteriophora* and *Heterorhabditis megidis*, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 671; *Steinernema feltiae*, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1115 and *Steinernema scapterisci*, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1116;
- (LIV) a preparation obtainable from *Bacillus subtilis*, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 72; or from a strain of *Bacillus thuringiensis* with the exception of compounds isolated from GC91 or from NCTC11821; The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 73;
- (LV) a preparation which contains insect-active fungi, preferably *Verticillium lecanii*, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1266; *Beauveria brogniartii*, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 85 and *Beauveria bassiana*, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 83;
- (LVI) a preparation which contains insect-active viruses, preferably *Neodiprion Sertifer* NPV, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 1342; *Mamestra brassicae* NPV, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 759 and *Cydia pomonella* granulosus virus, from The Pesticide Manual, 11<sup>th</sup>Ed. (1997), The British Crop Protection Council, London, page 291;

- (CLXXXI) 7-Chloro-2,3,4a,5-tetrahydro-2-[methoxycarbonyl(4-trifluoromethoxyphenyl)-carbamoyl]indole[1,2e]oxazolin-4a-carboxylate (DPX-MP062, Indoxycarb), from The Pesticide Manual, 11<sup>th</sup> Ed. (1997), The British Crop Protection Council, London, page 453;
- (CLXXXII) *N*-tert.-butyl-*N'*-(3,5-dimethylbenzoyl)-3-methoxy-2-methylbenzohydrazide (RH-2485, Methoxyfenozide), from The Pesticide Manual, 11<sup>th</sup> Ed. (1997), The British Crop Protection Council, London, page 1094; and
- (CLXXXIII) (*N'*-[4-methoxy-biphenyl-3-yl]-hydrazinecarboxylic acid isopropyl ester (D 2341), from Brighton Crop Protection Conference, 1996, 487-493;
- (R2) Book of Abstracts, 212th ACS National Meeting Orlando, FL, August 25-29 (1996), AGRO-020. Publisher: American Chemical Society, Washington, D.C. CONEN: 63BFAF.

As a consequence of the above details, a further essential aspect of the present invention relates to combination preparations for the control of parasites on warm-blooded animals, characterised in that they contain, in addition to a compound of formula I, at least one further active ingredient having the same or different sphere of activity and at least one physiologically acceptable carrier. The present invention is not restricted to two-fold combinations.

As a rule, the anthelmintic compositions according to the invention contain 0.1 to 99 % by weight, especially 0.1 to 95 % by weight of active ingredient of formula I, Ia or mixtures thereof, 99.9 to 1 % by weight, especially 99.8 to 5 % by weight of a solid or liquid admixture, including 0 to 25 % by weight, especially 0.1 to 25 % by weight of a surfactant.

The pour-on or spot-on method consists in applying the compound of formula I to a specific location of the skin or coat, advantageously to the neck or backbone of the animal. This takes place e.g. by applying a swab or spray of the pour-on or spot-on formulation to a relatively small area of the coat, from where the active substance is dispersed almost automatically over wide areas of the fur owing to the spreading nature of the components in the formulation and assisted by the animal's movements.

Pour-on or spot-on formulations suitably contain carriers, which promote rapid dispersement over the skin surface or in the coat of the host animal, and are generally regarded as spreading oils. Suitable carriers are e.g. oily solutions; alcoholic and isopropanolic solutions such as solutions of 2-octyldodecanol or oleyl alcohol; solutions in esters of monocarboxylic acids, such as isopropyl myristate, isopropyl palmitate, lauric acid oxalate, oleic acid oleyl

ester, oleic acid decyl ester, hexyl laurate, oleyl oleate, decyl oleate, capric acid esters of saturated fat alcohols of chain length  $C_{12}$ - $C_{18}$ ; solutions of esters of dicarboxylic acids, such as dibutyl phthalate, diisopropyl isophthalate, adipic acid diisopropyl ester, di-n-butyl adipate or also solutions of esters of aliphatic acids, e.g. glycols. It may be advantageous for a dispersing agent to be additionally present, such as one known from the pharmaceutical or cosmetic industry. Examples are 2-pyrrolidone, 2-(N-alkyl)pyrrolidone, acetone, polyethylene glycol and the ethers and esters thereof, propylene glycol or synthetic triglycerides.

The oily solutions include e.g. vegetable oils such as olive oil, groundnut oil, sesame oil, pine oil, linseed oil or castor oil. The vegetable oils may also be present in epoxidised form. Paraffins and silicone oils may also be used.

A pour-on or spot-on formulation generally contains 1 to 20 % by weight of a compound of formula I, 0.1 to 50 % by weight of dispersing agent and 45 to 98.9 % by weight of solvent.

The pour-on or spot-on method is especially advantageous for use on herd animals such as cattle, horses, sheep or pigs, in which it is difficult or time-consuming to treat all the animals orally or by injection. Because of its simplicity, this method can of course also be used for all other animals, including individual domestic animals or pets, and is greatly favoured by the keepers of the animals, as it can often be carried out without the specialist presence of the veterinarian.

Whereas it is preferred to formulate commercial products as concentrates, the end user will normally use dilute formulations.

Such compositions may also contain further additives, such as stabilisers, anti-foaming agents, viscosity regulators, binding agents or tackifiers, as well as other active ingredients, in order to achieve special effects.

Anthelmintic compositions of this type, which are used by the end user, similarly form a constituent of the present invention.

In each of the processes according to the invention for pest control or in each of the pest control compositions according to the invention, the active ingredients of formula I can be used in all of their steric configurations or in mixtures thereof.

The invention also includes a method of prophylactically protecting warm-blooded animals, especially productive livestock, domestic animals and pets, against parasitic helminths,

which is characterised in that the active ingredients of formula I or the active ingredient formulations prepared therefrom are administered to the animals as an additive to the feed, or to the drinks or also in solid or liquid form, orally or by injection or parenterally. The invention also includes the compounds of formula I according to the invention for usage in one of the said processes.

The following examples serve merely to illustrate the invention without restricting it, the term active ingredient representing a substance listed in table 1.

In particular, preferred formulations are made up as follows:

(% = percent by weight)

#### Formulation examples

##### 1. Emulsion concentrates

	a)	b)	c)
active ingredient	25 %	40 %	50 %
Ca dodecylbenzene sulphonate	5 %	8 %	6 %
castor oil polyethylene glycol ether (36 mols ethylene oxide)	5 %	-	-
tributylphenolpolyethylene glycol ether (30 mols ethylene oxide)	-	12 %	4 %
cyclohexanone	-	15 %	20 %
xylene mixture	65 %	25 %	20 %

From these concentrates, emulsions of any desired concentration may be prepared by diluting with water.

##### 2. Emulsion concentrates

	a)	b)	c)
active ingredient	10 %	8 %	60 %
octylphenol polyethylene glycol ether (4-5 mols ethylene oxide)	3 %	3 %	2 %
Ca dodecylbenzene sulphonate	3 %	4 %	4 %
castor oil polyethylene glycol ether (35 mols ethylene oxide)	4 %	5 %	4 %
cyclohexanone	30 %	40 %	15 %
xylene mixture	50 %	40 %	15 %

From these concentrates, emulsions of any desired concentration may be prepared by diluting with water.

### 3. Suspension concentrate

active ingredient	40 %
ethylene glycol	10 %
nonylphenol polyethylene glycol ether (15 mols ethylene oxide)	6 %
Na ligninsulphonate	10 %
carboxymethylcellulose	1 %
37% aqueous formaldehyde solution	0.2 %
silicone oil in the form of a 75% aqueous emulsion	0.8 %
water	32 %

The finely ground active ingredient is intimately mixed with the admixtures. In this way, a suspension concentrate is obtained, from which suspensions of any desired concentration can be prepared by diluting with water.

<u>4. Powder mixtures that are dispersible in water</u>	a)	b)	c)
active ingredient	25 %	50 %	75 %
Na ligninsulphonate	5 %	5 %	-
oleic acid	3 %	-	5 %
Na diisobutyl-naphthalene sulphonate	-	6 %	10 %
octylphenol polyethylene glycol ether (7-8 mols ethylene oxide)	-	2 %	-
highly dispersed silicic acid	5 %	10 %	10 %
kaolin	62 %	27 %	-

The active ingredient is mixed thoroughly with the admixtures and ground well in an appropriate mill. Wettable powders are obtained, which may be diluted with water to form suspensions of any desired concentration.

### 5. Dusts

	a)	b)
active ingredient	2 %	5 %
highly dispersed silicic acid	1 %	5 %
talc	97 %	-

kaolin - 90 %

By intimately mixing the carriers with the active ingredient and grinding the mixture, ready-to-use dusts are obtained.

#### 6. Granulates

	a)	b)
active ingredient	5 %	10 %
kaolin	94 %	-
highly dispersed silicic acid	1 %	-
attapulgit	-	90 %

The active ingredient is dissolved in methylene chloride, sprayed onto the carrier and the solvent subsequently concentrated by evaporation under vacuum. Granulates of this kind can be mixed with the fodder.

#### 7. Granulates

active ingredient	10 %
Na ligninsulphonate	2 %
carboxymethylcellulose	1 %
kaolin	87 %

The active ingredient is mixed with the admixtures, ground and moistened with water. This mixture is extruded and then dried in a stream of air.

#### 8. Granulates

active ingredient	3 %
polyethylene glycol (mw 200)	3 %
kaolin	94 %

(mw = molecular weight)

The finely ground active ingredient is evenly applied in a mixer to the kaolin which has been moistened with polyethylene glycol. In this way, dust-free coated granules are obtained.

#### 9. Tablets or boli

I	active ingredient	33.00 %
	methylcellulose	0.80 %
	silicic acid, highly dispersed	0.80 %
	corn starch	8.40 %
II	lactose, cryst.	22.50 %

corn starch	17.00 %
microcryst. cellulose	16.50 %
magnesium stearate	1.00 %

- I Methyl cellulose is stirred into water. After the material has swollen, silicic acid is stirred in and the mixture homogeneously suspended. The active ingredient and the corn starch are mixed. The aqueous suspension is worked into this mixture and kneaded to a dough. The resulting mass is granulated through a 12 M sieve and dried.
- II All 4 excipients are mixed thoroughly.
- III The preliminary mixes obtained according to I and II are mixed and pressed into tablets or boli.

#### 10. Injectables

##### A. Oily vehicle (slow release)

1. active ingredient 0.1-1.0 g  
groundnut oil ad 100 ml
2. active ingredient 0.1-1.0 g  
sesame oil ad 100 ml

Preparation: The active ingredient is dissolved in part of the oil whilst stirring and, if required, with gentle heating, then after cooling made up to the desired volume and sterile-filtered through a suitable membrane filter with a pore size of 0.22 mm.

##### B Water-miscible solvent (average rate of release)

- |   |           |
|---|-----------|
| active ingredient                               | 0.1-1.0 g |
| 4-hydroxymethyl-1,3-dioxolane (glycerol formal) | 40 g      |
| 1,2-propanediol                                 | ad 100 ml |
| an active ingredient                            | 0.1-1.0 g |
| glycerol dimethyl ketal                         | 40 g      |
| 1,2-propanediol                                 | ad 100 ml |

Preparation: The active ingredient is dissolved in part of the solvent whilst stirring, made up to the desired volume and sterile-filtered through a suitable membrane filter with a pore size of 0.22 mm.

##### C. Aqueous solubilisate (rapid release)

1. active ingredient	0.1-1.0 g
polyethoxylated castor oil (40 ethylene oxide units)	10 g
1,2-propanediol	20 g
benzyl alcohol	1 g
aqua ad inject.	ad 100 ml
2. active ingredient	0.1-1.0 g
polyethoxylated sorbitan monooleate (20 ethylene oxide units)	8 g
4-hydroxymethyl-1,3-dioxolane (glycerol formal)	20 g
benzyl alcohol	1 g
aqua ad inject.	ad 100 ml

Preparation: The active ingredient is dissolved in the solvents and the surfactant, and made up with water to the desired volume. Sterile filtration through an appropriate membrane filter of 0.22 mm pore size.

#### 11. Pour on

##### A.

active ingredient	5 g
isopropyl myristate	10 g
isopropanol	ad 100 ml

##### B

active ingredient	2 g
hexyl laurate	5 g
medium-chained triglyceride	15 g
ethanol	ad 100 ml

##### C.

active ingredient	2 g
oleyl oleate	5 g
N-methyl-pyrrolidone	40 g
isopropanol	ad 100 ml

The aqueous systems may also preferably be used for oral and/or intraruminal application.

The compositions may also contain further additives, such as stabilisers, e.g. where appropriate epoxidised vegetable oils (epoxidised coconut oil, rapeseed oil, or soybean oil);



antifoams, typically silicone oil; preservatives; viscosity regulators; binders; and tackifiers, as well as fertilisers or other chemical agents to achieve special effects.

Further biologically active substances or additives, which are neutral towards the compounds of formula I and do not have a harmful effect on the host animal to be treated, as well as mineral salts or vitamins, may also be added to the described compositions.

The following examples serve to illustrate the invention. They do not limit the invention. The letter 'h' stands for hour.


#### Preparation examples

##### Example 1: [N-(5-cyano-4-trifluoromethylthiazol-2-yl)]-3-phenylpropionamide

168 mg of 3-phenylpropionyl chloride are dissolved in 5 ml of dried dichloromethane, then 193 mg of 2-amino-5-cyano-4-trifluoromethylthiazole, followed by 0.46 ml of diisopropylethylamine and 23.6 mg of 4-dimethylaminopyridine, are added. The mixture is stirred for 20 h at room temperature in a nitrogen atmosphere, then washed with a little water, 25 ml of saturated sodium bicarbonate solution and finally 25 ml of saturated sodium chloride solution. After drying the organic phase with magnesium sulphate, filtering and concentrating by evaporation, the residue is purified in a HPLC column, yielding the title compound with a melting point of 156-159°C.

The substances named in the following tables 1 and 2 may also be prepared analogously to the above-described method. The values of the melting points are given in °C. Ph signifies phenyl.

Table 1



No.	X <sub>1</sub>	X <sub>2</sub>	R <sub>1</sub>	R <sub>3</sub> , R <sub>4</sub>	Q	R <sub>5</sub>	phys. data
1.1	N	N	Cl	H,H	-	Ph	
1.2	N	N	Cl	H,H	-	3-F-Ph	
1.3	N	N	Cl	H,H	-	3-Cl-Ph	
1.4	N	N	Cl	H,H	-	3-CH <sub>3</sub> O-Ph	
1.5	N	N	Cl	H,H	-	4-CF <sub>3</sub> -Ph	
1.6	N	N	Cl	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.7	N	N	Cl	H,H	-	2-Pyridyl	
1.8	N	N	Cl	H,H	-	3-Pyridyl	
1.9	N	N	Cl	H,H	-	2-Cl-3-pyridyl	
1.10	N	N	Cl	H,H	-	6-CH <sub>3</sub> O-3-pyridyl	
1.11	N	N	Cl	H,H	CH <sub>2</sub>	Ph	
1.12	N	N	Cl	H,H	CH <sub>2</sub>	3-F-Ph	
1.13	N	N	Cl	H,H	CH <sub>2</sub>	3-Cl-Ph	
1.14	N	N	Cl	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
1.15	N	N	Cl	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
1.16	N	N	Cl	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.17	N	N	Cl	H,H	CH <sub>2</sub>	2-Pyridyl	
1.18	N	N	Cl	H,H	CH <sub>2</sub>	3-Pyridyl	
1.19	N	N	Cl	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl	
1.20	N	N	Cl	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl	
1.21	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph	
1.22	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph	
1.23	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph	
1.24	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph	
1.25	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph	
1.26	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	

1.27	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl	
1.28	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl	
1.29	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl	
1.30	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl	
1.31	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph	
1.32	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph	
1.33	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph	
1.34	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
1.35	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
1.36	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.37	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl	
1.38	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl	
1.39	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl	
1.40	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl	
1.41	N	N	CCl <sub>3</sub>	H,H	-	Ph	
1.42	N	N	CCl <sub>3</sub>	H,H	-	3-F-Ph	
1.43	N	N	CCl <sub>3</sub>	H,H	-	3-Cl-Ph	
1.44	N	N	CCl <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph	
1.45	N	N	CCl <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph	
1.46	N	N	CCl <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.47	N	N	CCl <sub>3</sub>	H,H	-	2-Pyridyl	
1.48	N	N	CCl <sub>3</sub>	H,H	-	3-Pyridyl	
1.49	N	N	CCl <sub>3</sub>	H,H	-	2-Cl-3-pyridyl	
1.50	N	N	CCl <sub>3</sub>	H,H	-	6-CH <sub>3</sub> O-3-pyridyl	
1.51	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	Ph	m.p: 149-51°
1.52	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph	
1.53	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph	
1.54	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
1.55	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
1.56	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.57	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2-Pyridyl	
1.58	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Pyridyl	
1.59	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl	

1.60	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.61	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
1.62	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
1.63	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
1.64	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
1.65	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.66	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.67	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.68	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.69	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.70	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.71	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.72	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.73	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.74	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.75	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.76	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.77	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl
1.78	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl
1.79	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl
1.80	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.81	N	N	CF <sub>3</sub>	H,H	-	Ph
1.82	N	N	CF <sub>3</sub>	H,H	-	3-F-Ph
1.83	N	N	CF <sub>3</sub>	H,H	-	3-Cl-Ph
1.84	N	N	CF <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
1.85	N	N	CF <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
1.86	N	N	CF <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.87	N	N	CF <sub>3</sub>	H,H	-	2-Pyridyl
1.88	N	N	CF <sub>3</sub>	H,H	-	3-Pyridyl
1.89	N	N	CF <sub>3</sub>	H,H	-	2-Cl-3-pyridyl
1.90	N	N	CF <sub>3</sub>	H,H	-	6-CH <sub>3</sub> O-3-pyridyl
1.91	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
1.92	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph

1.93	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
1.94	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.95	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.96	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.97	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2-Pyridyl
1.98	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Pyridyl
1.99	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl
1.100	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.101	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
1.102	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
1.103	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
1.104	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
1.105	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.106	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.107	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.108	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.109	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.110	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.111	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.112	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.113	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.114	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.115	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.116	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.117	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl
1.118	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl
1.119	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl
1.120	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.121	N	C(CN)	Cl	H,H	-	Ph
1.122	N	C(CN)	Cl	H,H	-	3-F-Ph
1.123	N	C(CN)	Cl	H,H	-	3-Cl-Ph
1.124	N	C(CN)	Cl	H,H	-	3-CH <sub>3</sub> O-Ph
1.125	N	C(CN)	Cl	H,H	-	4-CF <sub>3</sub> -Ph

1.126	N	C(CN)	Cl	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.127	N	C(CN)	Cl	H,H	-	2-Pyridyl
1.128	N	C(CN)	Cl	H,H	-	3-Pyridyl
1.129	N	C(CN)	Cl	H,H	-	2-Cl-3-pyridyl
1.130	N	C(CN)	Cl	H,H	-	6-CH <sub>3</sub> O-3-pyridyl
1.131	N	C(CN)	Cl	H,H	CH <sub>2</sub>	Ph
1.132	N	C(CN)	Cl	H,H	CH <sub>2</sub>	3-F-Ph
1.133	N	C(CN)	Cl	H,H	CH <sub>2</sub>	3-Cl-Ph
1.134	N	C(CN)	Cl	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.135	N	C(CN)	Cl	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.136	N	C(CN)	Cl	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.137	N	C(CN)	Cl	H,H	CH <sub>2</sub>	2-Pyridyl
1.138	N	C(CN)	Cl	H,H	CH <sub>2</sub>	3-Pyridyl
1.139	N	C(CN)	Cl	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl
1.140	N	C(CN)	Cl	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.141	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
1.142	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
1.143	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
1.144	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
1.145	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.146	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.147	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.148	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.149	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.150	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.151	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.152	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.153	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.154	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.155	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.156	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.157	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl
1.158	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl

1.159	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl
1.160	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.161	N	C(CN)	CCl <sub>3</sub>	H,H	-	Ph
1.162	N	C(CN)	CCl <sub>3</sub>	H,H	-	3-F-Ph
1.163	N	C(CN)	CCl <sub>3</sub>	H,H	-	3-Cl-Ph
1.164	N	C(CN)	CCl <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
1.165	N	C(CN)	CCl <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
1.166	N	C(CN)	CCl <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.167	N	C(CN)	CCl <sub>3</sub>	H,H	-	2-Pyridyl
1.168	N	C(CN)	CCl <sub>3</sub>	H,H	-	3-Pyridyl
1.169	N	C(CN)	CCl <sub>3</sub>	H,H	-	2-Cl-3-pyridyl
1.170	N	C(CN)	CCl <sub>3</sub>	H,H	-	6-CH <sub>3</sub> O-3-pyridyl
1.171	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
1.172	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
1.173	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
1.174	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.175	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.176	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.177	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2-Pyridyl
1.178	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Pyridyl
1.179	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl
1.180	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.181	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
1.182	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
1.183	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
1.184	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
1.185	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.186	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.187	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.188	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.189	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.190	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.191	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph

1.192	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph	
1.193	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph	
1.194	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
1.195	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
1.196	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.197	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl	
1.198	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl	
1.199	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl	
1.200	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl	
1.201	N	C(CN)	CF <sub>3</sub>	H,H	-	Ph	
1.202	N	C(CN)	CF <sub>3</sub>	H,H	-	3-F-Ph	
1.203	N	C(CN)	CF <sub>3</sub>	H,H	-	3-Cl-Ph	
1.204	N	C(CN)	CF <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph	
1.205	N	C(CN)	CF <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph	
1.206	N	C(CN)	CF <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	m.p.: 133-6°
1.207	N	C(CN)	CF <sub>3</sub>	H,H	-	2-Pyridyl	
1.208	N	C(CN)	CF <sub>3</sub>	H,H	-	3-Pyridyl	
1.209	N	C(CN)	CF <sub>3</sub>	H,H	-	2-Cl-3-pyridyl	
1.210	N	C(CN)	CF <sub>3</sub>	H,H	-	6-CH <sub>3</sub> O-3-pyridyl	
1.211	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	Ph	m.p: 156-9°
1.212	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph	
1.213	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph	
1.214	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
1.215	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
1.216	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.217	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2-Pyridyl	
1.218	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Pyridyl	
1.219	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl	
1.220	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl	
1.221	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph	
1.222	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph	
1.223	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph	resin
1.224	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph	



1.225	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.226	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.227	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.228	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.229	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.230	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.231	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.232	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.233	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.234	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.235	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.236	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.237	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl
1.238	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl
1.239	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl
1.240	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.241	C(CN)	N	Cl	H,H	-	Ph
1.242	C(CN)	N	Cl	H,H	-	3-F-Ph
1.243	C(CN)	N	Cl	H,H	-	3-Cl-Ph
1.244	C(CN)	N	Cl	H,H	-	3-CH <sub>3</sub> O-Ph
1.245	C(CN)	N	Cl	H,H	-	4-CF <sub>3</sub> -Ph
1.246	C(CN)	N	Cl	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.247	C(CN)	N	Cl	H,H	-	2-Pyridyl
1.248	C(CN)	N	Cl	H,H	-	3-Pyridyl
1.249	C(CN)	N	Cl	H,H	-	2-Cl-3-pyridyl
1.250	C(CN)	N	Cl	H,H	-	6-CH <sub>3</sub> O-3-pyridyl
1.251	C(CN)	N	Cl	H,H	CH <sub>2</sub>	Ph
1.252	C(CN)	N	Cl	H,H	CH <sub>2</sub>	3-F-Ph
1.253	C(CN)	N	Cl	H,H	CH <sub>2</sub>	3-Cl-Ph
1.254	C(CN)	N	Cl	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.255	C(CN)	N	Cl	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.256	C(CN)	N	Cl	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.257	C(CN)	N	Cl	H,H	CH <sub>2</sub>	2-Pyridyl

1.258	C(CN)	N	Cl	H,H	CH <sub>2</sub>	3-Pyridyl
1.259	C(CN)	N	Cl	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl
1.260	C(CN)	N	Cl	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.261	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
1.262	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
1.263	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
1.264	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
1.265	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.266	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.267	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.268	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.269	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.270	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.271	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.272	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.273	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.274	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.275	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.276	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.277	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl
1.278	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl
1.279	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl
1.280	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.281	C(CN)	N	CCl <sub>3</sub>	H,H	-	Ph
1.282	C(CN)	N	CCl <sub>3</sub>	H,H	-	3-F-Ph
1.283	C(CN)	N	CCl <sub>3</sub>	H,H	-	3-Cl-Ph
1.284	C(CN)	N	CCl <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
1.285	C(CN)	N	CCl <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
1.286	C(CN)	N	CCl <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.287	C(CN)	N	CCl <sub>3</sub>	H,H	-	2-Pyridyl
1.288	C(CN)	N	CCl <sub>3</sub>	H,H	-	3-Pyridyl
1.289	C(CN)	N	CCl <sub>3</sub>	H,H	-	2-Cl-3-pyridyl
1.290	C(CN)	N	CCl <sub>3</sub>	H,H	-	6-CH <sub>3</sub> O-3-pyridyl

1.291	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
1.292	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
1.293	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
1.294	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.295	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.296	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.297	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2-Pyridyl
1.298	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Pyridyl
1.299	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl
1.300	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.301	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
1.302	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
1.303	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
1.304	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
1.305	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.306	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.307	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.308	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.309	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.310	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.311	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.312	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.313	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.314	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.315	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.316	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.317	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl
1.318	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl
1.319	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl
1.320	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.321	C(CN)	N	CF <sub>3</sub>	H,H	-	Ph
1.322	C(CN)	N	CF <sub>3</sub>	H,H	-	3-F-Ph
1.323	C(CN)	N	CF <sub>3</sub>	H,H	-	3-Cl-Ph

1.324	C(CN)	N	CF <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
1.325	C(CN)	N	CF <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
1.326	C(CN)	N	CF <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.327	C(CN)	N	CF <sub>3</sub>	H,H	-	2-Pyridyl
1.328	C(CN)	N	CF <sub>3</sub>	H,H	-	3-Pyridyl
1.329	C(CN)	N	CF <sub>3</sub>	H,H	-	2-Cl-3-pyridyl
1.330	C(CN)	N	CF <sub>3</sub>	H,H	-	6-CH <sub>3</sub> O-3-pyridyl
1.331	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
1.332	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
1.333	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
1.334	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.335	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.336	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.337	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2-Pyridyl
1.338	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Pyridyl
1.339	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl
1.340	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.341	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
1.342	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
1.343	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
1.344	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
1.345	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.346	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.347	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.348	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.349	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.350	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.351	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.352	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.353	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.354	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.355	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.356	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph

1.357	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl	
1.358	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl	
1.359	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl	
1.360	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl	
1.361	C(CN)	C(CN)	Cl	H,H	-	Ph	
1.362	C(CN)	C(CN)	Cl	H,H	-	3-F-Ph	
1.363	C(CN)	C(CN)	Cl	H,H	-	3-Cl-Ph	
1.364	C(CN)	C(CN)	Cl	H,H	-	3-CH <sub>3</sub> O-Ph	m.p: 195-200°
1.365	C(CN)	C(CN)	Cl	H,H	-	4-CF <sub>3</sub> -Ph	
1.366	C(CN)	C(CN)	Cl	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.367	C(CN)	C(CN)	Cl	H,H	-	2-Pyridyl	
1.368	C(CN)	C(CN)	Cl	H,H	-	3-Pyridyl	
1.369	C(CN)	C(CN)	Cl	H,H	-	2-Cl-3-pyridyl	
1.370	C(CN)	C(CN)	Cl	H,H	-	6-CH <sub>3</sub> O-3-pyridyl	
1.371	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	Ph	m.p: 246-8°
1.372	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	3-F-Ph	
1.373	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	3-Cl-Ph	
1.374	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
1.375	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
1.376	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.377	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	2-Pyridyl	
1.378	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	3-Pyridyl	
1.379	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl	
1.380	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl	
1.381	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph	
1.382	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph	
1.383	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph	
1.384	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph	
1.385	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph	
1.386	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
1.387	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl	
1.388	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl	
1.389	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl	

1.390	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.391	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.392	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.393	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.394	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.395	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.396	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.397	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl
1.398	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl
1.399	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl
1.400	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.401	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	Ph
1.402	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	3-F-Ph
1.403	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	3-Cl-Ph
1.404	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
1.405	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
1.406	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.407	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	2-Pyridyl
1.408	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	3-Pyridyl
1.409	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	2-Cl-3-pyridyl
1.410	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	6-CH <sub>3</sub> O-3-pyridyl
1.411	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
1.412	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
1.413	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
1.414	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.415	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.416	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.417	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2-Pyridyl
1.418	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Pyridyl
1.419	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl
1.420	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.421	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
1.422	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph


1.423	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
1.424	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
1.425	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.426	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.427	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.428	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.429	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.430	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.431	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.432	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.433	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.434	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.435	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.436	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.437	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl
1.438	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl
1.439	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl
1.440	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.441	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	Ph
1.442	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	3-F-Ph
1.443	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	3-Cl-Ph
1.444	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
1.445	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
1.446	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.447	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	2-Pyridyl
1.448	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	3-Pyridyl
1.449	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	2-Cl-3-pyridyl
1.450	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	6-CH <sub>3</sub> O-3-pyridyl
1.451	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
1.452	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
1.453	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
1.454	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.455	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph

1.456	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.457	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2-Pyridyl
1.458	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Pyridyl
1.459	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2-Cl-3-pyridyl
1.460	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl
1.461	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
1.462	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
1.463	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
1.464	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
1.465	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
1.466	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.467	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Pyridyl
1.468	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Pyridyl
1.469	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2-Cl-3-pyridyl
1.470	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	6-CH <sub>3</sub> O-3-pyridyl
1.471	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
1.472	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
1.473	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
1.474	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
1.475	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
1.476	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
1.477	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Pyridyl
1.478	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Pyridyl
1.479	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2-Cl-3-pyridyl
1.480	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	6-CH <sub>3</sub> O-3-pyridyl

---



Table 2



No.	X <sub>1</sub>	X <sub>2</sub>	R <sub>1</sub>	R <sub>3</sub> , R <sub>4</sub>	Q	R <sub>5</sub>	phys. data
2.1	N	N	Cl	H, H	-	Ph	
2.2	N	N	Cl	H, H	-	3-F-Ph	
2.3	N	N	Cl	H, H	-	3-Cl-Ph	
2.4	N	N	Cl	H, H	-	3-CH <sub>3</sub> O-Ph	
2.5	N	N	Cl	H, H	-	4-CF <sub>3</sub> -Ph	
2.6	N	N	Cl	H, H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.7	N	N	Cl	H, H	CH <sub>2</sub>	Ph	
2.8	N	N	Cl	H, H	CH <sub>2</sub>	3-F-Ph	
2.9	N	N	Cl	H, H	CH <sub>2</sub>	3-Cl-Ph	
2.10	N	N	Cl	H, H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
2.11	N	N	Cl	H, H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
2.12	N	N	Cl	H, H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.13	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph	
2.14	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph	
2.15	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph	
2.16	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph	
2.17	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph	
2.18	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.19	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph	
2.20	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph	
2.21	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph	
2.22	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
2.23	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
2.24	N	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.25	N	N	CCl <sub>3</sub>	H, H	-	Ph	

2.26	N	N	CCl <sub>3</sub>	H,H	-	3-F-Ph
2.27	N	N	CCl <sub>3</sub>	H,H	-	3-Cl-Ph
2.28	N	N	CCl <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
2.29	N	N	CCl <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
2.30	N	N	CCl <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.31	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
2.32	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
2.33	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
2.34	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.35	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.36	N	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.37	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
2.38	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.39	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.40	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
2.41	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.42	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.43	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
2.44	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.45	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.46	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.47	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.48	N	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.49	N	N	CF <sub>3</sub>	H,H	-	Ph
2.50	N	N	CF <sub>3</sub>	H,H	-	3-F-Ph
2.51	N	N	CF <sub>3</sub>	H,H	-	3-Cl-Ph
2.52	N	N	CF <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
2.53	N	N	CF <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
2.54	N	N	CF <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.55	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
2.56	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
2.57	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
2.58	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph

2.59	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.60	N	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.61	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
2.62	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.63	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.64	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
2.65	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.66	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.67	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
2.68	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.69	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.70	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.71	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.72	N	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.73	N	C(CN)	Cl	H,H	-	Ph
2.74	N	C(CN)	Cl	H,H	-	3-F-Ph
2.75	N	C(CN)	Cl	H,H	-	3-Cl-Ph
2.76	N	C(CN)	Cl	H,H	-	3-CH <sub>3</sub> O-Ph
2.77	N	C(CN)	Cl	H,H	-	4-CF <sub>3</sub> -Ph
2.78	N	C(CN)	Cl	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.79	N	C(CN)	Cl	H,H	CH <sub>2</sub>	Ph
2.80	N	C(CN)	Cl	H,H	CH <sub>2</sub>	3-F-Ph
2.81	N	C(CN)	Cl	H,H	CH <sub>2</sub>	3-Cl-Ph
2.82	N	C(CN)	Cl	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.83	N	C(CN)	Cl	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.84	N	C(CN)	Cl	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.85	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
2.86	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.87	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.88	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
2.89	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.90	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.91	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph

2.92	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.93	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.94	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.95	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.96	N	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.97	N	C(CN)	CCl <sub>3</sub>	H,H	-	Ph
2.98	N	C(CN)	CCl <sub>3</sub>	H,H	-	3-F-Ph
2.99	N	C(CN)	CCl <sub>3</sub>	H,H	-	3-Cl-Ph
2.100	N	C(CN)	CCl <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
2.101	N	C(CN)	CCl <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
2.102	N	C(CN)	CCl <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.103	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
2.104	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
2.105	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
2.106	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.107	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.108	N	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.109	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
2.110	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.111	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.112	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
2.113	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.114	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.115	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
2.116	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.117	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.118	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.119	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.120	N	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.121	N	C(CN)	CF <sub>3</sub>	H,H	-	Ph
2.122	N	C(CN)	CF <sub>3</sub>	H,H	-	3-F-Ph
2.123	N	C(CN)	CF <sub>3</sub>	H,H	-	3-Cl-Ph
2.124	N	C(CN)	CF <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph

2.125	N	C(CN)	CF <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph	
2.126	N	C(CN)	CF <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.127	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	Ph	m.p: 91-4°
2.128	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph	
2.129	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph	
2.130	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
2.131	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
2.132	N	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.133	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph	
2.134	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph	
2.135	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph	
2.136	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph	
2.137	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph	
2.138	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.139	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph	
2.140	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph	
2.141	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph	
2.142	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
2.143	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
2.144	N	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.145	C(CN)	N	Cl	H,H	-	Ph	
2.146	C(CN)	N	Cl	H,H	-	3-F-Ph	
2.147	C(CN)	N	Cl	H,H	-	3-Cl-Ph	
2.148	C(CN)	N	Cl	H,H	-	3-CH <sub>3</sub> O-Ph	
2.149	C(CN)	N	Cl	H,H	-	4-CF <sub>3</sub> -Ph	
2.150	C(CN)	N	Cl	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.151	C(CN)	N	Cl	H,H	CH <sub>2</sub>	Ph	
2.152	C(CN)	N	Cl	H,H	CH <sub>2</sub>	3-F-Ph	
2.153	C(CN)	N	Cl	H,H	CH <sub>2</sub>	3-Cl-Ph	
2.154	C(CN)	N	Cl	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph	
2.155	C(CN)	N	Cl	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph	
2.156	C(CN)	N	Cl	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph	
2.157	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph	

2.158	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.159	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.160	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
2.161	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.162	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.163	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
2.164	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.165	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.166	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.167	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.168	C(CN)	N	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.169	C(CN)	N	CCl <sub>3</sub>	H,H	-	Ph
2.170	C(CN)	N	CCl <sub>3</sub>	H,H	-	3-F-Ph
2.171	C(CN)	N	CCl <sub>3</sub>	H,H	-	3-Cl-Ph
2.172	C(CN)	N	CCl <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
2.173	C(CN)	N	CCl <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
2.174	C(CN)	N	CCl <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.175	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
2.176	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
2.177	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
2.178	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.179	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.180	C(CN)	N	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.181	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
2.182	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.183	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.184	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
2.185	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.186	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.187	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
2.188	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.189	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.190	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph

2.191	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.192	C(CN)	N	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.193	C(CN)	N	CF <sub>3</sub>	H,H	-	Ph
2.194	C(CN)	N	CF <sub>3</sub>	H,H	-	3-F-Ph
2.195	C(CN)	N	CF <sub>3</sub>	H,H	-	3-Cl-Ph
2.196	C(CN)	N	CF <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
2.197	C(CN)	N	CF <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
2.198	C(CN)	N	CF <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.199	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
2.200	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
2.201	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
2.202	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.203	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.204	C(CN)	N	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.205	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
2.206	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.207	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.208	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
2.209	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.210	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.211	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
2.212	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.213	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.214	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.215	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.216	C(CN)	N	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.217	C(CN)	C(CN)	Cl	H,H	-	Ph
2.218	C(CN)	C(CN)	Cl	H,H	-	3-F-Ph
2.219	C(CN)	C(CN)	Cl	H,H	-	3-Cl-Ph
2.220	C(CN)	C(CN)	Cl	H,H	-	3-CH <sub>3</sub> O-Ph
2.221	C(CN)	C(CN)	Cl	H,H	-	4-CF <sub>3</sub> -Ph
2.222	C(CN)	C(CN)	Cl	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.223	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	Ph

2.224	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	3-F-Ph
2.225	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	3-Cl-Ph
2.226	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.227	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.228	C(CN)	C(CN)	Cl	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.229	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
2.230	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.231	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.232	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
2.233	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.234	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.235	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
2.236	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.237	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.238	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.239	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.240	C(CN)	C(CN)	Cl	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.241	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	Ph
2.242	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	3-F-Ph
2.243	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	3-Cl-Ph
2.244	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
2.245	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
2.246	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.247	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
2.248	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
2.249	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
2.250	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.251	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.252	C(CN)	C(CN)	CCl <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.253	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
2.254	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.255	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.256	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph



2.257	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.258	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.259	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
2.260	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.261	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.262	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.263	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.264	C(CN)	C(CN)	CCl <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.265	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	Ph
2.266	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	3-F-Ph
2.267	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	3-Cl-Ph
2.268	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	3-CH <sub>3</sub> O-Ph
2.269	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	4-CF <sub>3</sub> -Ph
2.270	C(CN)	C(CN)	CF <sub>3</sub>	H,H	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.271	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	Ph
2.272	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-F-Ph
2.273	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-Cl-Ph
2.274	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.275	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.276	C(CN)	C(CN)	CF <sub>3</sub>	H,H	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.277	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	Ph
2.278	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-F-Ph
2.279	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-Cl-Ph
2.280	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	3-CH <sub>3</sub> O-Ph
2.281	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	4-CF <sub>3</sub> -Ph
2.282	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	-	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph
2.283	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	Ph
2.284	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-F-Ph
2.285	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-Cl-Ph
2.286	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	3-CH <sub>3</sub> O-Ph
2.287	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	4-CF <sub>3</sub> -Ph
2.288	C(CN)	C(CN)	CF <sub>3</sub>	-CH <sub>2</sub> CH <sub>2</sub> -	CH <sub>2</sub>	2,5-(CH <sub>3</sub> O) <sub>2</sub> -Ph

---

### Biological Examples:

#### 1. In-vivo test on *Trichostrongylus colubriformis* and *Haemonchus contortus* on Mongolian gerbils (*Meriones unguiculatus*) using subcutaneous injection

Six to eight week old Mongolian gerbils are infected by artificial feeding with ca. 2000 third instar larvae each of *T. colubriformis* and *H. contortus*. 6 days after infection, the gerbils are lightly anaesthetised with N<sub>2</sub>O and treated by subcutaneous injection into the neck area with the test compounds, dissolved in a mixture of 2 parts DMSO and 1 part polyethylene glycol 400, in quantities of 100, 32 and 10 - 0.1 mg/kg. On day 9 (3 days after treatment), when most of the *H. contortus* that are still present are 4th instar larvae and most of the *T. colubriformis* are immature adults, the gerbils are killed in order to count the worms. The efficacy is calculated as the % reduction of the number of worms in each gerbil, compared with the geometric average of number of worms from 8 infected and untreated gerbils.

In this test, a vast reduction in nematode infestation is achieved with compounds of formula I.

Totally analogous results are obtained upon oral administration of the active ingredient.

#### 2. Insecticidal stomach toxicant activity on *Spodoptera littoralis*

Potted cotton plants at the 5-leaf stage are each sprayed with an acetonic/aqueous test solution containing 1, 3, 12.5 or 50 ppm of the compound to be tested.

After drying of the spray deposit, the plants are colonised with ca. 30 larvae (L<sub>1</sub> stage) of *Spodoptera littoralis*. Two plants are used per test compound and per test species. The test is carried out at ca. 24°C and at 60% relative humidity. Evaluations and intermediate evaluations on moribund animals, larvae and feeding damage are made after 24, 48 and 72 h.

The compounds of formula I achieve total mortality after 24 h at a concentration of active ingredient of only 3 ppm.

#### 3. Activity on phytotoxic acarids

##### OP-sensitive *Tetranychus urticae*

The primary leaves of bean plants (*Phaseolus vulgaris*) are covered 16 hours before the test with a mass-cultivated piece of leaf infested with *T. urticae*. After removing the piece of leaf, the plants that are infested with all stages of the mites are sprayed to drip point with a test solution containing either 0.2, 0.4 or 1.6 ppm of the compound to be tested. The

temperature in the greenhouse is ca. 25°C. After 7 days, an evaluation of the percentage of mobile stages (adults and nymphs) and of eggs is made under a microscope.

The compounds of formula I achieve total mortality at a concentration of active ingredient of 0.4 ppm.

#### 4. Activity on L<sub>1</sub> larvae of *Lucilia sericata*

1 ml of an aqueous suspension of the active substance to be tested is admixed with 3 ml of a special larvae growth medium at ca. 50°C, so that a homogenate of either 250 or 125 ppm of active ingredient content is obtained. Ca. 30 *Lucilia* larvae (L<sub>1</sub>) are used in each test tube sample. After 4 days, the mortality rate is determined. The compounds of formula I attain 100% activity with 250 ppm.

#### 5. Acaricidal activity on *Boophilus microplus* (Biarra strain)

A piece of sticky tape is attached horizontally to a PVC sheet, so that 10 fully engorged female ticks of *Boophilus microplus* (Biarra strain) can be adhered thereto by their backs, side by side, in a row. Using an injection needle, 1 µl of a liquid is injected into each tick. The liquid is a 1:1 mixture of polyethylene glycol and acetone and it contains, dissolved therein, a certain amount of active ingredient chosen from 1, 0.1 or 0.01 µg per tick. Control animals are given an injection without active ingredient. After treatment, the animals are kept under normal conditions in an insectarium at ca. 28°C and at 80% relative humidity until oviposition takes place and the larvae have hatched from the eggs of the control animals. The activity of a tested substance is determined by IR<sub>90</sub>, i.e. an evaluation is made of the dosage of active ingredient at which 9 out of 10 female ticks (=90%) lay eggs that are infertile even after 30 days.

The compounds of formula I attain an IR<sub>90</sub> of 0.1 µg.

#### 6. In vitro efficacy on engorged female *Boophilus microplus* (BIARRA):

4x10 engorged female ticks of the OP-resistant BIARRA strain are adhered to a sticky strip and covered for 1 hour with a cotton-wool ball soaked in an emulsion or suspension of the test compound in concentrations of 500, 125, 31 and 8 ppm respectively. Evaluation takes place 28 days later based on mortality, oviposition and hatched larvae.

An indication of the activity of the test compounds is shown by the number of females that

- die quickly before laying eggs,
- survive for some time without laying eggs,

- lay eggs in which no embryos are formed,
  - lay eggs in which embryos form, from which no larvae hatch, and
  - lay eggs in which embryos form, from which larvae normally hatch within 26 to 27 days.
- In this test, the compounds of formula I effect more than 80% rapid mortality of the female ticks.

#### 7. Contact action on *Aphis craccivora*

Pea seedlings that have been infested with all stages of development of the aphids are sprayed with a solution of active ingredient prepared from an emulsion concentrate, the solution containing 50, 25 or 12.5 ppm of active ingredient, as desired. After 3 days, an evaluation is made of more than 80% of aphids that are either dead or have fallen off. Only at this level of activity is a preparation classified as effective.

The compounds of formula I achieve total mortality (= 100%) at a concentration of 12.5 ppm.

#### 8. Larvicidal activity on *Aedes aegypti*

A sufficient quantity of a 0.1% acetic solution of the active ingredient for a chosen concentration of 10, 3.3 or 1.6 ppm to be attained, is added by pipette to the surface of 150 ml of water in a container. After evaporation of the acetone, the container is covered with ca. 30-40 3-day old *Aedes* larvae. After 1, 2 and 5 days, the mortality is tested.

In this test, the compounds of formula I at a concentration of 1.6 ppm effect complete mortality of all larvae after only one day.

#### 9. *In vivo* efficacy on adult *Ctenocephalides felis* on domestic cats after oral treatment

The test substances are given orally to domestic cats in a gelatin capsule before or after feeding, the dose varying between 0.5 and 20 mg/kg. On days 1, 3, 7 and 10 after treatment, each cat is exposed to 100 fleas (ca. 50 male and ca. 50 female), depending on the result of previous flea colonisation. The efficacy (in % reduction in flea numbers) is based on the number of living fleas found after combing for 10 minutes one day after each new flea colonisation, whereby the efficacy in % corresponds to the arithmetic average of the number of living fleas on control animals minus the number of living fleas on the treated animals, divided by the arithmetic average of the number of living fleas on control animals and multiplied by 100.

The dying fleas found in the cat cages and by combing are collected, placed in an incubator at 28°C and 70% relative humidity and after 24 hours are tested for survival/mortality. If the majority of dying fleas die, the test compound is regarded as a flea adulticide, and if the majority survive, the test compound shows "knock-down" activity.

In this test, the compounds of formula I effect at least 80% mortality of the fleas.

10. In vivo efficacy on adult *Ctenocephalides felis* on domestic cats after spot-on treatment

The test substances are given to domestic cats as spot-on treatment, the dose varying between 0.5 and 10 mg/kg. On days 1, 3, 7 and 10 after treatment, each cat is exposed to 100 fleas (ca. 50 male and ca. 50 female), depending on the result of previous flea colonisation.

The efficacy (in % reduction in flea numbers) is based on the number of living fleas found after combing for 10 minutes one day after each new flea colonisation, whereby the efficacy in % corresponds to the arithmetic average of the number of living fleas on control animals minus the number of living fleas on the treated animals, divided by the arithmetic average of the number of living fleas on control animals and multiplied by 100.

The dying fleas found in the cat cages and by combing are collected, placed in an incubator at 28°C and 70% relative humidity and after 24 hours are tested for survival/mortality. If the majority of dying fleas die, the test compound is regarded as a flea adulticide, and if the majority survive, the test compound shows "knock-down" activity.

In this test, the compounds of formula I effect more than 90% mortality of the fleas after 35 days.

11. In vitro efficacy on nymphs of *Amblyomma hebraeum*

About 5 fasting nymphs are placed in a polystyrene test tube containing 2 ml of the test compound in solution, suspension or emulsion.

After immersion for 10 minutes, and shaking for 2x10 seconds on a vortex mixer, the test tubes are blocked up with a tight wad of cotton wool and rotated. As soon as all the liquid has been soaked up by the cotton wool ball, it is pushed half-way into the test tube which is still being rotated, so that most of the liquid is squeezed out of the cotton-wool ball and flows into a Petri dish below.

The test tubes are then kept at room temperature in a room with daylight until evaluated. After 14 days, the test tubes are immersed in a beaker of boiling water. If the ticks begin to move in reaction to the heat, the test substance is inactive at the tested concentration, otherwise the ticks are regarded as dead and the test substances regarded as active at the tested concentration. All substances are tested in a concentration range of 0.1 to 100 ppm. In this test, the compounds of formula I effect more than 80% mortality of the ticks.

12. Activity against *Dermanyssus gallinae*

2 to 3 ml of a solution containing 10 ppm active ingredient, and ca. 200 mites (*Dermanyssus gallinae*) at different stages of development are added to a glass container which is open at the top. Then the container is closed with a wad of cotton wool, shaken for 10 minutes until the mites are completely wet, and then inverted briefly so that the remaining test solution can be absorbed by the cotton wool. After 3 days, the mortality of the mites is determined by counting the dead individuals and indicated as a percentage.

The compounds of formula I show good activity against *Dermanyssus gallinae*.


13. Activity against *Musca domestica*

A sugar cube is treated with a solution of the test substance in such a way that the concentration of test substance in the sugar, after drying over night, is 250 ppm. The cube treated in this way is placed on an aluminium dish with wet cotton wool and 10 adult *Musca domestica* of an OP-resistant strain, covered with a beaker and incubated at 25°C. The mortality rate is determined after 24 hours.

In this test, the compounds of formula I show good activity against *Musca domestica*.

What we claim is:

## 1. Compounds of formula



wherein

$R_1$  is halogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -haloalkoxy,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -haloalkylthio,  $C_1$ - $C_6$ -alkoxy- $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkylthio- $C_1$ - $C_6$ -alkyl,  $C_3$ - $C_8$ -cycloalkyl or phenyl;

$R_2$  is hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy- $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkylsulphonyl,  $C_3$ - $C_8$ -cycloalkyl,  $-COOR_6$ ,  $-CONR_7R_8$ ,  $-COR_6$ , allyl,  $-CH_2-O-R_6$ ; or heteroaryl, arylcarbonyl, arylsulphonyl or aryl- $C_1$ - $C_6$ -alkyl which are each either unsubstituted or substituted once or many times by substituents selected from the group comprising  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy, halogen, cyano, hydroxy, amino and nitro, whereby if the number of substituents is greater than 1, the substituents may be identical or different;

$R_3$  and  $R_4$ , independently of one another, are hydrogen,  $C_1$ - $C_6$ -alkyl or together with the C-atom to which they are bonded, a  $C_3$ - $C_7$ -cycloalkyl ring;

$Q$  is a single bond,  $C_1$ - $C_6$ -alkylene,  $C_2$ - $C_6$ -alkenylene,  $C_2$ - $C_6$ -alkynylene or  $C_1$ - $C_6$ -alkylenoxy;

$R_5$  is aryl or heterocyclyl, each of which is either unsubstituted or substituted once or many times by substituents selected from the group comprising  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy, aryloxy, halogen, cyano, hydroxy, amino and nitro, whereby if the number of substituents is greater than 1, the substituents may be identical or different.

$R_6$  is  $C_1$ - $C_6$ -alkyl, phenyl or benzyl;


$R_7$  and  $R_8$  independently of one another, are hydrogen or  $C_1$ - $C_6$ -alkyl;

$X_1$  and  $X_2$  independently of one another, are N, C(CN),  $-C(COOR_6)=$ ,  $-C(COR_6)=$ ,  $-C(SOR_6)=$ ,  $-C(CONR_7R_8)=$  or  $-C(NO_2)=$ ; and


$X_3$  is O or S;

whereby  $X_1$  is other than C(CN) if  $R_5$  is phenyl that is either unsubstituted or substituted once to many times.

2. Process for the preparation of compounds of formula I according to claim 1, whereby a compound of formula



which is known or may be produced analogously to corresponding known compounds, and wherein  $R_1$ ,  $R_2$ ,  $X_1$  and  $X_2$  are defined as given for formula I, is reacted with a compound of formula



which is known or may be produced analogously to corresponding known compounds, in which  $X_3$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $Q$  are defined as for formula I, and  $Z$  is a leaving group, optionally in the presence of a basic catalyst, and if desired, a compound of formula I which is obtainable by this process or in another way, or an enantiomer thereof, may be converted into another compound of formula I or an enantiomer thereof, a mixture of enantiomers which is obtainable by this process is separated and the desired enantiomer isolated.

3. Composition for the control of pests, which contains as active ingredient at least one compound of formula I according to claim 1, in addition to carriers and/or dispersants.
4. Use of compounds of formula I according to claim 1 in the control of pests.
5. Method of controlling pests, whereby a pesticidally active amount of at least one compound of formula I according to claim 1 is used on the pests or on the locus thereof.
6. Use of a compound of formula I according to claim 1 in a process for controlling parasites on warm-blooded animals.



7. Use of a compound of formula I according to claim 1 in the preparation of a pharmaceutical composition against parasites.



(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number  
WO 01/46165 A3

(51) International Patent Classification: C07D 277/56,  
285/08, 275/02, 333/38, 417/12, 409/12, A61K 31/426,  
31/433, 31/4439, 31/425, 31/381, A61P 33/00, A01N  
43/78, 43/836, 43/80, 43/10, 43/40

(FR). BOUVIER, Jacques [CH/CH]; Rue des Parcs 2,  
CH-2000 Neuchatel (CH).

(21) International Application Number: PCT/EP00/12751

(74) Agent: BECKER, Konrad; Novartis AG, Corporate In-  
tellectual Property, Patent & Trademark Dept., CH-4002  
Basel (CH).

(22) International Filing Date:  
14 December 2000 (14.12.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
2305/99 16 December 1999 (16.12.1999) CH

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except AT, US*): NO-  
VARTIS AG [CH/CH]; Lichtstrasse 35, CH-4056 Basel  
(CH).

Published:  
— with international search report

(71) Applicant (*for AT only*): NOVARTIS-ERFINDUNGEN  
VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT];  
Brunner Strasse 59 A-1230 Vienna (AT).

(88) Date of publication of the international search report:  
6 December 2001


(72) Inventors; and

(75) Inventors/Applicants (*for US only*): DUCRAY, Pierre  
[FR/FR]; 15A, rue de Huningue, F-68300 Saint-Louis

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

WO 01/46165 A3

(54) Title: N-HETEROARYL-AMIDES AND THEIR USE AS PARASITICIDES



(57) Abstract: The invention relates to compounds of general formula (I), wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub> and Q have the significances given in claim 1, and optionally the enantiomers thereof. The active ingredients have advantageous pesticidal properties. They are especially suitable for the control of pests on domestic animals and livestock.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/12751

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D277/56 C07D285/08 C07D275/02 C07D333/38 C07D417/12  
C07D409/12 A61K31/426 A61K31/433 A61K31/4439 A61K31/425  
A61K31/381 A61P33/00 A01N43/78 A01N43/836 A01N43/80

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A01N A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 26251 A (BAYER AKTIENGESELLSCHAFT) 24 July 1997 (1997-07-24) the whole document ---	1-7
X	WO 97 18198 A (BAYER AKTIENGESELLSCHAFT) 22 May 1997 (1997-05-22) cited in the application the whole document ---	1-7
X	EP 0 566 138 A (HODOGAYA CHEMICAL CO., LTD.) 20 October 1993 (1993-10-20) the whole document ---	1-7
X	FR 2 014 527 A (E. MERCK AKTIENGESELLSCHAFT) 17 April 1970 (1970-04-17) the whole document ---	1-7
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

19 June 2001

Date of mailing of the international search report

06/07/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Allard, M

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/12751

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01N43/10 A01N43/40

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 455 356 A (LILLY INDUSTRIES LIMITED) 6 November 1991 (1991-11-06) page 8, line 56 ---	1-3
X	GB 1 551 735 A (LILLY INDUSTRIES LIMITED) 30 August 1979 (1979-08-30) examples 26, 74 and 75 ---	1-3
X	OHKUBO M ET AL: "Studies on cerebral protective agents. VIII. Synthesis of 2-aminothiazoles and 2-thiazolecarboxamides with anti-anoxic activity" CHEMICAL & PHARMACEUTICAL BULLETIN, vol. 43, no. 9, September 1995 (1995-09), pages 1497-1504, XP002169988 page 1499, table I, compounds 2i and 2k --- -/--	1-3

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

19 June 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Allard, M

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/12751

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 64, no. 13, 20 June 1966 (1966-06-20) Columbus, Ohio, US; abstract no. 19589f, SHARMA S C: "Synthesis of new local anesthetics. VIII. Synthesis of 4,5-disubstituted thiazoles" XP002169989 abstract & INDIAN J. CHEM., vol. 4, no. 1, 1966, pages 33-36, ---	1-3
X	US 4 675 404 A (BERNARDI L ET AL) 23 June 1987 (1987-06-23) examples 33 and 34 ---	1-3
P,X	WO 00 06566 A (ZENECA LIMITED) 10 February 2000 (2000-02-10) the whole document ---	1-7
E	WO 01 36415 A (NOVARTIS AG ET AL) 25 May 2001 (2001-05-25) the whole document -----	1-7

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-7 (all partly)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claims may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons it appears impossible to execute a meaningful search and/or to issue a complete search report over the whole breadth of the above mentioned claims.

The search and the report for those claims can only be considered complete for compounds of formula I according to claim 1 wherein X1 and X2 are N or C(CN)

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

I. International Application No

PCT/EP 00/12751

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9726251 A	24-07-1997	DE 19601139 A	17-07-1997
		AU 712985 B	18-11-1999
		AU 1438797 A	11-08-1997
		BR 9706986 A	20-07-1999
		CA 2242905 A	24-07-1997
		CN 1208408 A	17-02-1999
		EP 0877741 A	18-11-1998
		HU 9900971 A	28-07-1999
		JP 2000503307 T	21-03-2000
		US 5942528 A	24-08-1999
WO 9718198 A	22-05-1997	DE 19542372 A	15-05-1997
		AU 7563396 A	05-06-1997
		BR 9611475 A	13-07-1999
		CN 1202162 A	16-12-1998
		EP 0861241 A	02-09-1998
		JP 2000502046 T	22-02-2000
		US 5972843 A	26-10-1999
		US 6008366 A	28-12-1999
EP 566138 A	20-10-1993	DE 69322118 D	24-12-1998
		DE 69322118 T	20-05-1999
		JP 6025199 A	01-02-1994
FR 2014527 A	17-04-1970	DE 1770819 A	13-01-1972
		DE 1924830 A	19-11-1970
		AT 287707 B	10-02-1971
		BE 735655 A	05-01-1970
		CH 547819 A	11-04-1974
		DK 124207 B	25-09-1972
		ES 369148 A	16-06-1971
		GB 1226913 A	31-03-1971
		IL 32487 A	16-05-1974
		NL 6910070 A	08-01-1970
		SE 359300 B	27-08-1973
EP 455356 A	06-11-1991	CA 2039955 A	11-10-1991
		JP 4224570 A	13-08-1992
GB 1551735 A	12-09-1979	AR 217240 A	14-03-1980
		AT 346843 B	27-11-1978
		AT 412276 A	15-04-1978
		AU 503150 B	23-08-1979
		AU 1461976 A	08-12-1977
		BE 842579 A	03-12-1976
		BG 27359 A	12-10-1979
		CA 1085860 A	16-09-1980
		CH 616933 A	30-04-1980
		CH 612427 A	31-07-1979
		CS 200494 B	15-09-1980
		DD 125075 A	30-03-1977
		DE 2625285 A	23-12-1976
		DK 243276 A	06-12-1976
		EG 12226 A	31-12-1978
		ES 448590 A	16-11-1977
		FI 761585 A,B,	06-12-1976
		FR 2313049 A	31-12-1976
		GR 60380 A	23-05-1978



# INTERNATIONAL SEARCH REPORT

Information on patent family members

i. International Application No

PCT/EP 00/12751

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1551735 A		HU 174191 B	28-11-1979
		IE 43474 B	11-03-1981
		IL 49692 A	31-07-1980
		JP 51146468 A	16-12-1976
		LU 75070 A	18-01-1978
		MX 3483 E	16-12-1980
		NL 7606178 A	07-12-1976
		NO 761913 A,B,	07-12-1976
		NZ 181011 A	13-11-1978
		OA 5346 A	28-02-1981
		PH 16077 A	20-06-1983
		PT 65182 A,B	01-07-1976
		SE 428561 B	11-07-1983
		SE 7606226 A	06-12-1976
		SU 639451 A	25-12-1978
		US 4163048 A	31-07-1979
		US 4152442 A	01-05-1979
		US 4158659 A	19-06-1979
		US 4166818 A	04-09-1979
		YU 137176 A	31-10-1982
		ZA 7603209 A	25-01-1978
US 4675404 A	23-06-1987	AT 13766 T	15-06-1985
		AU 553809 B	31-07-1986
		AU 8610582 A	27-01-1983
		BE 893872 A	16-11-1982
		CA 1191840 A	13-08-1985
		CH 652407 A	15-11-1985
		DE 3264121 D	18-07-1985
		DK 324182 A	22-01-1983
		EP 0070562 A	26-01-1983
		FI 822528 A,B,	22-01-1983
		FR 2510115 A	28-01-1983
		IT 1210480 B	14-09-1989
		JP 58024584 A	14-02-1983
		ZA 8205141 A	27-04-1983
WO 0006566 A	10-02-2000	AU 5055099 A	21-02-2000
		EP 1100798 A	23-05-2001
WO 0136415 A	25-05-2001	NONE	

**THIS PAGE BLANK (USPTO)**